



REVIEW

Recent advances in 4-hydroxycoumarin chemistry. Part 2: Scaffolds for heterocycle molecular diversity



Moaz M. Abdou ^{a,d,*}, Rasha A. El-Saeed ^b, Samir Bondock ^{b,c}

^a Egyptian Petroleum Research Institute, Nasr City, P.O. 11727, Cairo, Egypt

^b Department of Chemistry, Faculty of Science, Mansoura University, ET-35516 Mansoura, Egypt

^c Department of Chemistry, Faculty of Science, King Khalid University, 9004 Abha, Saudi Arabia

^d The Institute of Scientific and Industrial Research, Osaka University, 8-1 Mihogaoka, Ibaraki-shi, Osaka 567-0047, Japan

Received 5 December 2014; accepted 26 June 2015

Available online 3 July 2015

KEYWORDS

4-Hydroxycoumarin;
Annulated heterocycles;
Multicomponent reactions;
Green chemistry

Abstract The present paper aims to review the synthetic applicability of 4-hydroxycoumarin in heterocyclic synthesis during the period from 1996 up to now. This compound can be used as building blocks for five, six, and seven-membered heterocycles as well as fused rings.

© 2015 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	975
2. Synthesis of monocyclic heterocyclic compounds	975
2.1. Synthesis of five-membered systems with two heteroatoms	975
2.1.1. Pyrazolin-5-ones.	975
2.1.2. Imidazol-2(3H)-ones.	975
2.1.3. 1,2-Benzisoxazoles	976
2.2. Synthesis of five-membered ring systems with more than two heteroatoms	976
2.2.1. Dithiazoles	976
2.3. Synthesis of six-membered heterocycles containing one heteroatom	976
2.3.1. Quinoline	976
2.4. Synthesis of seven-membered heterocycles containing two heteroatoms	976
2.4.1. 1,4-Diazepines	976
3. Synthesis of fused heterocyclic compounds	976

* Corresponding author. Tel.: +20 1000409279.

E-mail address: moaz.chem@gmail.com (M.M. Abdou).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

3.1.	[5-6] Ring system	976
3.1.1.	Pyrano[4,3-d]oxazol-4-ones	976
3.1.2.	Oxathiol[5,4-c]-2H-chromen-4-ones	976
3.1.3.	Dihydrofuran and furocoumarins	977
3.2.	Fused [6-6]ring system	982
3.2.1.	Benzopyrano[2,3-b]pyridine	982
3.2.2.	Pyrido[3,2-c]coumarins	982
3.2.3.	Chromeno[3,4-b][4,7]phenanthroline	983
3.2.4.	Chromeno[4,3-b]benzo[f]quinolin-6-one	983
3.2.5.	Pyrano benzopyrans	983
3.2.6.	3,4-Dihydro-2 <i>H</i> ,5 <i>H</i> -1-benzopyrano[4,3-b]pyran-5-one	992
3.2.8.	Benzopyrano[4,3-d]pyrimidines	995
3.2.9.	Benzopyranobenzothiazinones	997
3.2.10.	Chromeno[3,4-e][1,3]oxazine	997
3.2.11.	Pyrano[3,4-e][1,4,3]oxathiazines	997
3.3.	[7-6] Ring system	997
3.3.1.	Benzopyrano[1,3]diazepines	997
4.	Conclusion	997
	References	998

1. Introduction

4-Hydroxycoumarin is a well-known versatile synthon and important reagents in the realm of heterocyclic synthesis (Abdolmohammadi, 2014; Abdou et al., 2013; Aggarwal et al., 2013; Ahadi et al., 2014; Altieri et al., 2010; Azizi et al., 2014; Bakthadoss and Sivakumar, 2014; Chen et al., 2013; Darehkordi et al., 2014; Davarpanah and Kiasat, 2014; Eskandari et al., 2013; Ghosh and Khan, 2014; Golzar et al., 2013; Gu et al., 2009; Hossain et al., 2013; Hossaini et al., 2013; Jashari et al., 2014; Jin et al., 2013; Jung and Park, 2009; Lee et al., 2012; Mei et al., 2013; Metwally et al., 2013, 2012a, 2012b, 2012c, 2012d; Mungra et al., 2011; Nair et al., 2014; Nishino et al., 2014; Pal et al., 2013; Ponpandian and Muthusubramanian, 2014; Rad-Moghadam et al., 2014; Rajawat et al., 2014; Saha, 2013; Sato et al., 2013; Siddiqui and Khan, 2013, 2014; Xiao et al., 2014; Zhao and Du, 2014; Ziarani and Hajiabbasi, 2013). It is of particular interest as a very promising reagent for cascade heterocyclization, which will undoubtedly become one of the main approaches to the targeted synthesis of heterocycles in the near future, in the rapidly-rising field of combinatorial chemistry.

In a continuation of our research program aiming at the utilization of cyclic 1,3-diketone compounds as scaffolds for heterocycle molecular diversity (Abdou, 2017a, 2017b, 2017, 2018, 2017, 2013b, 2013c; Abdou et al., 2013, 2012), the first part of this review article (Abdou, 2019) is dealt with most recent advances in 4-hydroxycoumarin chemistry made since 1996 up to date and concentrates on synthesis, chemical reactivity and reactions of 4-hydroxycoumarin. The second part intends to illustrate the research efforts that occurred in the application of 4-hydroxycoumarin in heterocyclic synthesis covering the same period of time. The presence of reactive centers in this compound provides ample opportunities to synthesize a great variety of novel compounds under relatively mild conditions and using simple laboratory equipments. Thus, the two parts are complementary and display current trends in 4-hydroxycoumarin chemistry.

In this literature survey, the reactions involving 4-hydroxycoumarin occur with high regioselectivity and its

course can easily be controlled by changing reaction conditions and varying substituents in the molecules of initial compounds. The heterocyclic compounds are obtained in a single step with high yield and they are reported in order of the increase of (i) the number of rings, (ii) the size of such rings and (iii) the number of heteroatoms present. The sequence of heteroatoms followed is: nitrogen, oxygen and sulfur. The site of fusion in fused heterocycles is indicated by the numbers and letters and the numbering of the heterocyclic ring systems is that reported by chemical abstracts.

2. Synthesis of monocyclic heterocyclic compounds

2.1. Synthesis of five-membered systems with two heteroatoms

2.1.1. Pyrazolin-5-ones


Froggett et al. (1997) described the reaction of 4-hydroxycoumarin **1** with a 1.5 equimolar excess of 4-chlorophenylhydrazine **2** in toluene under reflux, with azeotropic removal of water, furnished the 1-(4-chlorophenyl)-3-(2-hydroxy-phenyl)-2-pyrazolin-5-one **3** (Scheme 1).

The solvent-free reactions of substituted hydrazine hydrochlorides **4** with **1** in the presence of triethylamine were carried out by heating in the absence of solvent (90 min, 90–100 °C). In all cases, 1-aryl-3-(2-hydroxyphenyl)pyrazol-5-ones **6a-c** were formed along with the corresponding 4-(arylhydrazino) coumarins **5a-c** (Strakova et al., 2009) (Scheme 2).


3-(2-Hydroxyphenyl)-1-methyl-1*H*-pyrazol-5-one **8** was obtained from the reaction of **1** with methylhydrazine **7** in ethanol. The reaction mixture was refluxed under N₂ flow (Zhao et al., 2010) (Scheme 3).

2.1.2. Imidazol-2(3*H*)-ones

The one-pot condensation of **1** with arylglyoxals **9** and ureas **10** in refluxing ethanol in the presence of a catalytic amount of acetic acid gave a colorless crystalline products that were identified as 4-aryl-5-(4-hydroxy-2-oxo-2*H*-chromen-3-yl)-1*H*-imidazol-2(3*H*)-ones **11** (Kolos et al., 2009) (Scheme 4).




Scheme 1



4, 5, 6	a	b	c
Ar	Ph	4-Cl-C ₆ H ₄	4-Br-C ₆ H ₄
Yield % (5/6)	51/18	53/9	52/15

Scheme 2



Scheme 3

2.1.3. 1,2-Benzisoxazoles

Reaction of **1** with hydroxylamine hydrochloride in refluxing methanol represents one of the most successful strategies to attain 1,2-benzisoxazole-3-acetic acid **12**, which has been widely exploited because of their known important biological activities and application in different therapies (Lamani et al., 2009) (Scheme 5).

2.2. Synthesis of five-membered ring systems with more than two heteroatoms

2.2.1. Dithiazoles

The reaction of **1** with 4,5-dichloro-5*H*-1,2,3-dithiazolium chloride **13** in the presence of pyridine in dichloromethane at room temperature gave the corresponding 5-alkylidene-5*H*-4-chloro-1,2,3-dithiazoles **14** as a single stereoisomer in 87% yield (Jeon and Kim, 1999) (Scheme 6).

2.3. Synthesis of six-membered heterocycles containing one heteroatom

2.3.1. Quinoline

In 2010, a fast and solvent-free method was described for the synthesis of substituted quinoline derivatives *via* Friedländer synthesis is promoted, either under conventional

heating or under MW irradiation, catalyzed by a derivatized silica bearing alkylsulfonic acid groups. The reaction of **1** with 2-aminobenzophenone **15** afforded 7-phenyl-6*H*-chromeno[4,3-*b*]quinolin-6-one **16** and 2-(4-phenylquinolin-2-yl)phenol **17**. The addition of activated molecular-sieve powder to the reacting mixture increased the yield of **16** (40%) while dramatically reducing the yield of **17** (11%), *o*-hydroxy acetophenone **18** was the main product (about 50%) generated by the partial degradation of excess 4-hydroxycoumarin (Garella et al., 2010) (Scheme 7).

2.4. Synthesis of seven-membered heterocycles containing two heteroatoms

2.4.1. 1,4-Diazepines

It was reported that the condensation of **1** with *o*-phenylenediamine **19** in refluxing toluene (Hamdi et al., 2006) or under MW irradiation (Kidwai et al., 2005) afforded 4-(2-hydroxyphenyl)-2,3-dihydro-1*H*-1,5-benzodiazepin-2-one (Scheme 8).

3. Synthesis of fused heterocyclic compounds


3.1. [5-6] Ring system

3.1.1. Pyrano[4,3-*d*]oxazol-4-ones

Ray and Paul (2004) published the one-pot synthesis of benzo[1]pyrano[4,3-*d*]oxazol-4-one **22** consisting of the condensation of **1** with formamide **21** under reflux (Scheme 9).

3.1.2. Oxathiol[5,4-*c*]-2*H*-chromen-4-ones

Reaction of 3-(chlorothio)-3-fluoro-2-(trifluoromethyl)-2-propanoic acid methyl ester **23** with **1** leads to sulfonylation accompanied by cyclization giving rise to fused 2-(2,2,2-trifluoro-1-




	9,10	Ar	R ₁	R ₂	Yield%
a	—	Ph	H	H	70
b	—	4-Me-Ph	H	H	68
c	—	4-Cl-Ph	H	H	69
d	—	4-Br-Ph	H	H	63
e	—	4-I-Ph	H	H	63
f	—	Ph	Ph	H	69
g	—	4-Me-Ph	Ph	H	55
h	—	4-Cl-Ph	Ph	H	58
i	—	Ph	Me	Me	47
j	—	4-Me-Ph	Me	Me	45
k	—	4-Cl-Ph	Me	Me	45

Scheme 4


methoxycarbonylethylidene)-oxathio[4,5-*c*]-2*H*-chromen-4-one **24** (Kovregin et al., 2001) (Scheme 10).

3.1.3. Dihydrofuran and furocoumarins

Furocoumarins are an important class of heterocyclic compounds possessing anticoagulant, insecticide, anthelmintic, hypnotic, antifungal, and HIV protease inhibition activities (Baichurin et al., 2013; Karami et al., 2013; Khan et al., 2013).



Scheme 5



Scheme 6

3.1.3.1. Oxidative cycloaddition reaction mediated by metal salts. The oxidative addition reaction of carbon-centered radicals to alkenes mediated by metal salts Ag (I), Ce (IV), Yb (III), Ru (II), and Mg(II) has received considerable attention over the last decade in organic synthesis for construction of carbon–carbon bonds.


3.1.3.1.1. Using Ag (I). Lee and Kim (1997) have reported a facile and simple method for the synthesis of dihydrofurans, by the addition of silver (I)/Celite to the mixture of **1** and with ethyl vinyl ether **25** in acetonitrile under reflux (Scheme 11).

In a similar manner, medium- and large-sized ring substituted furans **28** could be also achieved via cycloaddition of **1** with a variety of vinyl sulfides **27** in refluxing acetonitrile (Lee and Kim, 1997) (Scheme 12).

3.1.3.1.2. Using Ce(IV). An efficient method for one-step synthesis of substituted 2-arylfurans has been developed by ceric (IV) ammonium nitrate (CAN) mediated oxidative cycloaddition of **1** with phenylacetylene **29** at 0 °C in acetonitrile giving linear and angular furocoumarin derivatives **30** and **31** as a mixture of regioisomers in good yields (Lee et al., 1998) (Scheme 13).


Lee et al. (2000) has also found that CAN(IV) was the much superior reagent for this oxidative cycloaddition than Mn(OAc)₃·2H₂O and Ag₂CO₃/Celite. The reaction was typically carried out at 80 °C starting from **1** with methyl methacrylate **32** in the presence of CAN and excess amounts of sodium bicarbonate in acetonitrile giving the sole biologically interesting dihydrofurocoumarin **33** in 35% yield (Scheme 14).

Appendino et al. (1998) showed that treatment of **1** with 3-buten-2-ol and cerium (IV) ammonium nitrate (CAN) in




	Condition	time /min.	Yield % (16/17/18)
a	Conventional heating	60	13/20/---
b	80 °C, MW irradiation	5	22/36/---
c	80 °C, MW irradiation, Molecular sieve	5	40/11/50

Scheme 7




	Condition	Yield %
a	PhMe , reflux	80
b	MW	89

Scheme 8



Scheme 9




Scheme 11


acetonitrile at room temperature give a mixture of diastereomers **35** and **36** (Scheme 15). No substantial improvement of the ratio between angular and linear adducts could be achieved by changing solvent or metal oxidant and varying the temperature (Scheme 15).

It was also reported that the reaction of **1** with a range of alkenes in acetonitrile containing cerium (IV) ammonium


nitrate (CAN) afforded the corresponding furo[3,2-c]pyranones **38** and furo[2,3-b]pyranone derivatives **39**. It should be noted that the reactions of **1** with vinyl benzoate and methylstyrene produced exclusively the furo[3,2-c]benzopyranone derivatives, as the sole isolated product; no more than a trace amount of the corresponding furo[2,3-b]benzopyranone derivatives was obtained in each of these




Scheme 10



Scheme 12



Scheme 13



Scheme 14

reactions, while formation of both the furopyranone derivatives was observed using other alkenes (Kobayashi et al., 1999) (Scheme 16).

There has been a considerable interest in the use of CAN oxidation reactions in ionic liquids. Hence, reaction of **1** with methylstyrene **40** and CAN in 1-*n*-butyl-3-methylimidazolium tetrafluoroborate [bmim][BF₄]-dichloromethane(1:9), gave tricycle **41** in 70% yield after 3 h. The CAN reaction in [bmim][BF₄]-dichloromethane (40 °C) is therefore more efficient and/or faster than reactions (at r.t. or at 40 °C) in acetonitrile. One further and important advantage to the use of [bmim][BF₄] is that on workup, the cerium byproducts form a precipitate at the end of the reaction (Bar et al., 2003) (Scheme 17).


3.1.3.1.3. *Ruthenium(II)*. Cadierno et al. (2008) developed a simple strategy for the synthesis of multi-substituted

furocoumarins by using a catalytic system consisting of the 16-electron allyl-ruthenium(II) complex [Ru(η³-2-C₃H₄Me)(CO)(dppf)][SbF₆](dppf = 1,1'-bis(diphenylphosphino)ferrocene) and trifluoroacetic acid (TFA) has been used to promote the coupling between secondary propargylic alcohol **42** and **1** to give 3-(4-methoxyphenyl)-2-methyl-furo[3,2-*c*]chromen-4-one **43** (Scheme 18).


3.1.3.1.4. *Manganese(III) acetate*. Yilmaz et al. (2008) have obtained 2,3-dihydro-4*H*-furo[3,2-*c*]chromen-4-ones **45** from the radical cyclization of **1** mediated by manganese(III) acetate with electron rich alkenes **44** in 36–86% yields (Scheme 19).

3.1.3.1.5. *Multicopper oxidases (Laccases)*. Laccase (*Agaricus bisporus*)-catalyzed domino reaction of **1** with catechols **46** using atmospheric oxygen as the oxidant delivers for the synthesis of 10-substituted 8,9-dihydroxy-6*H*-benzo furo[3,2-*c*]chromen-6-ones **47** as single regioisomers with yields of 61–99% (Leutbecher et al., 2011, 2005) (Scheme 20). Some of these compounds have been made accessible by crude peroxidase from onion solid waste (Angeleska et al., 2013).

3.1.3.2. *[4 + 1] Cycloaddition reaction followed by a [1,3]-H shift*. A [1 + 4] approach was developed by Majumdar et al. to prepare 3-hydroxy-2,3-dihydrofuro[3,2-*c*][1]benzopyran-4-one **49** through the reaction of **1** with chloroacetaldehyde **48** in the presence of aq. potassium carbonate at room temperature for 1.5 h (Majumdar and Bhattacharyya, 1997) (Scheme 21).




Scheme 15




37,38,39	R ₁	R ₂	R ₃	Yield% (38/39)
a	CH ₃	CH ₃	CH ₃	25/45
b	H	(CH ₂) ₃	(CH ₂) ₃	31/31
c	OAc	H	H	17/8
d	OBz	H	H	15/---
e	Ph	CH ₃	H	62/---

Scheme 16



Scheme 17



44,45	R ₁	R ₂	R ₃	Yield %
a	Ph	H	Me	50
b	Me	Ph	H	81
c	Ph	Ph	Et	86
d	Ph	C ₃ H ₇	Ph	36
e	Me	OCOCH ₃	H	54

Scheme 19

In a similar manner, Bondock et al. (2011) reported that antimicrobial active 2-(5-hydroxy-3-methyl-1-phenyl-1*H*-pyrazole-4-yl)-4-*H*-furo[3,2-*c*]chromen-4-one was obtained via treatment of 2-chloro-1-(5-hydroxy-3-methyl-1-phenyl-1*H*-pyrazol-4-yl)ethanone **50** with **1** in sodium ethoxide solution (Scheme 22).

2,3-Dihydro-2-hydroxy-3-(4-hydroxy-2-oxo-2*H*-chromen-3-yl)furo[3,2-*c*]chromen-4-one **53** could be successfully prepared through the reaction of **1** with a 40% aqueous solution of glyoxal **52** in acetonitrile under reflux (Lamani et al., 2009) (Scheme 23).


Interaction of **1** with bromonitroalkenes **54** in aqueous sodium acetate and tetrabutylammonium bromide (TBAB) gave 2,3-dihydrofuro[3,2-*c*]chromen-4-ones **55** (Xie et al., 2011) (Scheme 24).

3.1.3.3. Multicomponent reactions.


3.1.3.3.1. Using of conventional synthesis. Several reports on the synthesis of furocoumarins via the multicomponent reactions of **1** with various arylaldehydes **56** and isocyanide **57** have been published during the last decade. In most of the

cases, the yields of the desired furocoumarins **58** were moderate. Moreover, extended reaction times of up to 24 h under reflux conditions in toxic solvents such as benzene or toluene (Nair et al., 2002; Panja et al., 2012; Zhu et al., 2011) were required. These drawbacks restrictedly limit the applications for furocoumarin library construction (Scheme 25).

Consequently, there is need to develop a simple, rapid, eco-friendly and easy experimental and products' isolation procedure for the synthesis of these compounds.




Scheme 18



46,47	R	t/h	Yield %
a	H	7	85
b	Me	3	99
c	OMe	5	61
d	CO ₂ Me	4	89

Scheme 20




Scheme 21


3.1.3.3.2. Using of microwave irradiation. Microwaves accelerated the synthesis of a range of furo[3,2-c]chromen-4-ones **59** via a three-component reaction of **1**, arylaldehyde **56** and cyclohexyl isocyanide **57** (Shaabani and Teimouri, 2003; Wu, 2006). By applying a microwave protocol (DMF, 150 °C, 5 min), reduction in reaction times (24 h → 3–5 min), higher yields and less by-products were observed compared to the conventional synthesis (Scheme 26).

3.1.3.3.3. Using of water. In fact, choice of solvent is one of the problems to face in order to perform eco-efficient processes. The presence of water was proven to be determinate for the success achieved by Shaabani et al. (2004). Further reasons that make water unique compared to other organic solvents are that it is cheap, not inflammable, and more important, it is not toxic. The environment-friendly one-pot three-component condensation reaction of **1**, *p*-substituted benzaldehyde **56**, and alkyl or aryl isocyanides **57** in water afforded furocoumarins **59** in good yields (Shaabani et al., 2004) (Scheme 27).

3.1.3.3.4. Using of solvent-free synthesis. Adib et al. (2009) have devoted considerable attention to an efficient and direct solvent-free synthesis of bioactive furo[3,2-c]chromenes via a



Scheme 23




54,55	Ar	Yield%
a	4-MeO-C ₆ H ₄	93
b	4-Me-C ₆ H ₄	87
c	Ph	90
e	4-Cl-C ₆ H ₄	96
f	4-Br-C ₆ H ₄	83

Scheme 24


one-pot and solvent-free reaction. Thus a mixture of **1**, aldehyde **56**, and isocyanide **57** underwent a one-pot addition reaction under an argon atmosphere and solvent-free conditions. The reaction proceeded at 180 °C and was complete within 1.5 h to produce *N*-alkyl-3-arylfuro[3,2-c]chromenes **59** in 91–95% yields (Scheme 28).

3.1.3.3.5. Using combination of solvent-free supported reagents and microwave. Shaabani et al. (2005b) have shown the combination of solvent-free supported reagents and microwave in the absence of solid supports and classical heating. One-pot method simply involves microwave irradiation of a mixture of *p*-substituted benzaldehyde **59** and **1** with isocyanides **60** in the presence of montmorillonite K10 to afford the corresponding products **61**. The reaction is completed in all cases within 4–5 min. Also, in this approach the use of large volume of benzene or toluene is avoided, work-up considerably simplified, and safety is increased by reducing the risk of over pressure and explosions (Scheme 29).

3.1.3.4. Using of electrochemical routes. One of the most successful strategies for the synthesis of 6*H*-benzofuro[3,2-c][1]ben




Scheme 22



56,58	Ar	t / h	Yield%
a	2-NO ₂ -Ph	24	69
b	4-NO ₂ -Ph	16	93
c	4-Cl-Ph	24	61
e	-Ph	24	77
f	4-MeO-Ph	16	62
g	4-Me-Ph	24	72

Scheme 25



56,57,59	Ar	R	Yield%
a	4-NO ₂ -Ph	<i>t</i> -Bu	84
b	4-NO ₂ -Ph	cyclohexyl	94
c	4-NO ₂ -Ph	2, 6-(CH ₃) ₂ -Ph	92
d	4-Cl-Ph	cyclohexyl	85
e	4-MeO-Ph	cyclohexyl	90

Scheme 26

56,57,59	Ar	R	Yield%
a	4-NO ₂ -Ph	<i>t</i> -Bu	79
b	4-NO ₂ -Ph	<i>c</i> -Hex	85
c	4-NO ₂ -Ph	2, 6-(CH ₃) ₂ -Ph	91
d	4-Cl-Ph	<i>c</i> -Hex	78
e	4-MeO-Ph	<i>c</i> -Hex	71

Scheme 27

zopyran-6-one derivatives **61** is electrochemical routes. Nematollahi et al. demonstrated the electrochemical oxidation of catechol (**60a**), 3-methylcatechol (**60b**), 3-methoxycatechol (**60c**), 2,3-dihydroxybenzoic acid (**60d**) and 3,4-dihydroxybenzoic acid (**60f**) with **1** as a nucleophile in the presence of potassium ferricyanide as an oxidizing agent in aqueous solution (Nematollahi et al., 2005; Golabi and Nematollahi, 1997a, 1997b) (Scheme 30).

3.2. Fused [6-6]ring system

3.2.1. Benzopyrano[2,3-*b*]pyridine


Bandyopadhyay et al. (2010) reported the one-pot synthesis of the 1-benzopyrano[2,3-*b*]pyridine moiety **63** via Knoevenagel condensation by heating an equimolar mixture of **1** and

2-(monosubstituted amino)chromone-3-carbaldehyde **62** in boiling ethanol containing a catalytic amount of pyridine (Scheme 31).

Maiti et al. (2010) described the one-pot reaction of 2-(*N*-cinnamyl-*N*-aryl)amino-4-oxo-4*H*-1-benzopyran-3-carbaldehydes **64** with **1** by heating in ethanol in the presence of a catalytic amount of pyridine afforded substituted 11-allyl-6,11-dihydro-6-(4-hydroxy-2-oxo-2*H*-1-benzopyran-3-yl)-1-benzopyrano[2,3-*b*]quinolin-5*H*-5-one **65** (Scheme 32).


3.2.2. Pyrido[3,2-*c*]coumarins

The work of Pandya et al. (2006) demonstrated that various diarylpyrido[3,2-*c*]coumarins **67** were synthesized in one step by reacting **1** with α,β -unsaturated ketones **66** in the presence of ammonium acetate and acetic acid (Scheme 33).




56,57,59	Ar	R	Yield%
a	Ph	c-Hex	91
b	Ph	t-Bu	95
c	4-MePh	t-Bu	95

Scheme 28



56,57,59	Ar	R	Yield%
a	4-NO ₂ -Ph	t-Bu	72
b	4-NO ₂ -Ph	c-Hex	80
c	4-NO ₂ -Ph	2, 6-(CH ₃) ₂ -Ph	70
d	4-Cl-Ph	c-Hex	72
e	4-MeO-Ph	c-Hex	74

Scheme 29



60,61	R ₁	R ₂	R ₃	Yield %
a	H	H	H	96
b	CH ₃	H	H	95
c	OCH ₃	H	H	98
d	H	CO ₂ H	H	90
e	H	H	CO ₂ H	90

Scheme 30

3.2.3. Chromeno[3,4-b][4,7]phenanthroline

A short and environmental-friendly synthesis of chromeno[3,4-b][4,7]phenanthroline derivatives **70** was accomplished by three-component reactions involving an aromatic aldehyde **68**, 6-aminoquinoline **69** and **1** in water, under microwave irradiation without the use of any catalyst (Zhuang et al., 2008) (Scheme 34).

3.2.4. Chromeno[4,3-b]benzo[f]quinolin-6-one

Chromeno[4,3-b]benzo[f]quinolin-6-one derivatives **72** were obtained in good to high yields via the reaction of *N*-arylidene-naphthalen-2-amine **71** with **1** in aqueous media catalyzed by triethylbenzyl ammonium chloride (TEBAC) (Wang et al., 2006, 2005) (Scheme 35).


3.2.5. Pyrano benzopyrans

Pyrano[3,2-c]chromen-5-one (pyrano[3,2-c]coumarin) is the core of important natural products and heterocyclic structures. Molecules with such a nucleus exhibit a wide range of biological and pharmacological properties, such as antioxidant, anti-cancer, anti-inflammatory, antiviral, and antibacterial activities (Ahadi et al., 2013; Banard et al., 2002; Ghandi et al., 2013; Magiatis et al., 1998; Mahdavinia and Peikarporsan, 2013; Mail et al., 2002; Shafiee et al., 2011; Siddiqui, 2014). Many naturally occurring compounds with pyrano[3,2-c]coumarin skeletons have been isolated, such as isoethuliacoumarin A, isoethuliacoumarin B, isoethuliacoumarin C, ethuliacoumarin A, ethuliacoumarin B, and pterophyllin. In addition, pyrano[3,2-c]coumarin derivatives are also used as novel photochromers with promising applications in many photonic materials (Huang et al., 2007).

3.2.5.1. 2H,5H-pyrano[3,2-c]benzo[b]pyran-2,5-dione. Many methods for the synthesis of 2H, 5H-pyrano[3,2-c]benzo[b]pyran-2,5-dione derivatives have been reported successively.


3.2.5.1.1. Using alkyl-2-substituted-3-dimethyl-amino-propenoates. Jakse et al. (2004) observed that the treatment of ethyl (2E)-3-dimethylamino-2-(1H-indol-3-yl)-propenoate **73** with **1** in acetic acid under reflux afforded 3-(1H-indol-3-yl)-2H,5H-pyrano[3,2-c]chromene-2,5-dione **74** (Scheme 36).

There are several reports in the literatures (Jukic et al., 2001a, 2001b; Sorsak et al., 1998; Toplak et al., 1997) about




62,63	R ₁	R ₂	t / h	Yield%
a	Me	Ph	5	85
b	H	Ph	6	79
c	H	Et	5.5	79

Scheme 31



64,65	R ₁	R ₂	t / h	Yield%
a	H	4-MeC ₆ H ₄	5	80
b	Me	Ph	4.5	76


Scheme 32



R₁, R₂, R₃ = H, OCH₃


R₄ = H, 4-MeOPh

Scheme 33




68,70	a	b	c	d	e	f	g	h	i	j
R	4-FC ₆ H ₄	4-ClC ₆ H ₄	4-BrC ₆ H ₄	4-CH ₃ OC ₆ H ₄	4-NO ₂ C ₆ H ₄	3-NO ₂ C ₆ H ₄	4-CH ₃ C ₆ H ₄	C ₆ H ₅	2,4-Cl ₂ C ₆ H ₃	thien-2-yl
Conditions (time, yield)	(6min,93%)	(6min,93%)	(5min,94%)	(6min,95%)	(5min,94%)	(5min,95%)	(5min,95%)	(5min,94%)	(6min,93%)	(5min,95%)


Scheme 34



Scheme 35




Scheme 36




75,76	R	R ₁	R ₂	Yield%
a	Me	CO ₂ Et	Pyr	30
b	Et	CO ₂ Et	Pyr	30
c	Me	CN	Pyr	47
d	Et	CN	Pyr	58
e	CO ₂ Et	CO ₂ Ph	CO ₂ Ph	93
f	CO ₂ Et	CO ₂ Et	CO ₂ Et	41

Scheme 37




Scheme 38




79,80	R ₁	X	t/h	Yield%
a	CO ₂ Me	O	2	81
b	CO ₂ Me	N-COPh	2	93
c	CH ₂ CO ₂ Ph	O	3	37

Scheme 39



Scheme 40



Scheme 41

85,86	R	t/min (A / B)	Yield % (A/ B)
a	Me	2.5/3	66/81
b	Ph	144/48	64/85
c	4-Br-Ph	96/48	65/ 83
d	5-Br-Ph	168/---	53/---
e	3,5-Br ₂ -Ph	24/48	80/41
f	2-MeO-5-Br-Ph	---/144	---/ 76


A : R,R-DBFOX/Ph+ Ni(ClO₄)₂·6H₂O + TMP
B : R,R-DBFOX/Ph+ Ni(ClO₄)₂·6H₂O + TMP+ Ac₂O

Scheme 42

the reaction of alkyl 2-[(2,2-disubstituted) ethenyl]amino-3-(dimethylamino)propenoate **75** with **1** in acetic acid for preparation of corresponding 2*H*,5*H*-pyrano[3,2-*c*]benzo[*b*]pyran-2,5-dione derivatives **76** (Scheme 37).

In 1999, it has been reported that the preparation of 3-(benzyloxycarbonyl)amino-2*H*,5*H*-pyrano[3,2-*c*]benzopyran-2,5-dione **78** was accomplished by the treatment of methyl-2-(benzyloxy carbonyl)amino-3-dimethylaminopropenoate **77** with **1** under reflux (Toplak et al., 1999) (Scheme 38).

3.2.5.1.2. Using substituted-tetrahydrofuran-2-one or oxopyrrolidine. A unique one step “ring switching” synthesis of (*S*)-3-(2,5-dioxo-2*H*,5*H*-benzo[*b*]pyrano[4,3,-*b*]pyran-3-yl)ester **80**



87,88	R₁	R₂	R₃	Yield % (Condition)
a	CN	Me	Me	60 (A)
b	CN	Naph.	H	85 (A)
c	NO ₂	2-MeOC ₆ H ₄	H	80 (B)
d	NO ₂	3,4-MeOC ₆ H ₄	H	80 (B)
e	NO ₂	4-MeOC ₆ H ₄	H	85 (B)
f	CO ₂ Me	4-MeOC ₆ H ₄	H	66 (C)
g	CN	4-MeOC ₆ H ₄	H	52 (C)

A : Morpholine /EtOH
 B : TEA /EtOH
 C : NaOMe/MeOH


Scheme 43

is notable when **79** is reacted with **1** in boiling acetic acid (Mihelic et al., 2001) (Scheme 39).


An elegant, efficient one-pot synthesis of 3-hydroxyethyl-2-pyrano[3,2-c]benzopyran-2,5(6*H*)-dione **82** was achieved by the condensation of **1** with the sodium salt of α -formyl- γ -butyrolactone **81** in the presence of ammonium acetate (Toche et al., 1999) (Scheme 40).

3.2.5.1.3. Using Pechmann-Duisberg reaction. The Pechmann-Duisberg reaction was employed by Abd El-Aziz et al. (2007) to synthesis 3-(chloromethyl)-2*H*-benzo[*h*]chromen-2-one **84** via condensation of **1** with ethyl 4-chloroacetoacetate **83** in the presence of nitrobenzene and anhydrous aluminum chloride (Scheme 41).

3.2.5.1.4. Using double catalytic activation conditions. Japanese research group presented an effective enantioselective synthetic method based on a concept of double catalytic activation by the use of catalytic amounts of chiral Lewis acid



Scheme 45




93,94	R₁	R₂	Time/h (A/B)	Yield% (A/B)
a	3-NO ₂	CN	10 / 11	88 / 91
b	4-F	CN	10 / 8	93 / 93
c	3,4-(CH ₃ O) ₂	CN	10 / 10	89 / 84
d	4-CH ₃	CN	8 / 10	73 / 80
e	H	CO ₂ Et	8 / 8	90 / 85
f	4-F	CO ₂ Et	6 / 7	89 / 90
g	4-CH ₃	CO ₂ Et	7 / 11	93 / 70
h	4-Cl	CO ₂ Et	7 / 7	97 / 80


A : H₂O, TEBA, 90 °C.
 B : KF-montmorillonite, DMF.

Scheme 46

and amine base, respectively. Thus, under the reaction conditions using 2,2,6,6-tetramethylpiperidine (TMP) and the enantio pure complex catalyst derived from the *R,R*-DBFOX/Ph ligand and nickel(II) perchlorate hexahydrate, the reaction of




Scheme 44



56,95,96	R	R ₁	Yield %
a	Me	CN	92
b	Me(CH ₂) ₁₀	CN	90
c	Me(CH ₂) ₅	CN	89
d	EtCH(Me)	CN	87
e	Me(CH ₂) ₈	CN	88
f	CH(CH ₃) ₂	NH ₂ -CS	74

Scheme 47



56,96	R	Yield %
a	i-Pr	80
b	PhCH ₂	77
c	cyclohexyl	66
d	heptyl	59
e	cyclohex-3-enyl	90
f	i-Bu	87
g	Ph(CH ₂) ₂	61

Scheme 48

1 with 4-bromo-1-crotonoyl-3,5-dimethylpyrazole **85** in tetrahydrofuran at room temperature in the presence of acetic anhydride (Itoh and Kanemasa, 2003) or the absence of it (Itoh et al., 2005), to give the corresponding enol lactones in a good yield (Scheme 42).

3.2.5.2. Synthesis of amino-substituted pyranopyranones. Amino-substituted pyranopyranones are useful intermediates for the synthesis of bioactive natural products and pharmaceutical drugs (Dekamin et al., 2013; Jaggavarapu et al., 2014;

Khan et al., 2014, 2011; Mehrabi and Abusaidi, 2010; Niknam and Jamali, 2012; Pansuriya et al., 2009; Prasanna and Raju, 2011; Shaabani et al., 2007; Shaterian and Oveisi, 2011; Tabatabaieian et al., 2012; Wang et al., 2010; Zheng and Li, 2011).

3.2.5.2.1. Two-component condensation. A number of publications have been taken out for Michael reactions **1** with a variety of substituted cinnamionitrile **87** in the presence of bases such as morpholine (Dyachenko and Rusanov, 2006; Kislyi et al., 1999), triethylamine (Mahmoud et al., 2009), sodium methoxide (Nesterov et al., 2005) resulted in the corresponding 2-amino-4-aryl-4H,5H-pyrano[3,2-c]chromenes **88** (Scheme 43).


In a similar fashion, it was noted that Michael cycloaddition reaction of **1** with various substituted α -cyanocinnamionitriles **89** in ethanolic piperidine under reflux afforded pyrano[3,2-c]coumarins **90** (Bedair et al., 2000; El-Agrody et al., 2000; Abd El-Wahab et al., 2011) (Scheme 44).

Shaabani et al. (2008) described an isocyanide-catalyzed reaction between tetracyanoethylene **91** and **1** to afford 2-amino-5-oxopyrano[3,2-c]chromene-3,4,4(5H)-tricarbonitrile **92** in high yield at room temperature (Scheme 45).

The conversion of substituted cinnamionitriles **93** and **1** into 2-amino-4-aryl-4H,5H-pyrano[3,2-c][1]benzopyran-5-one derivatives **94** can be efficiently performed in water as a solvent using a catalytic amount of triethyl-benzylammonium chloride (TEBA) (Wang et al., 2004) or KF-montmorillonite (Tu et al., 2004) (Scheme 46).


3.2.5.2.2. Three component condensation.

3.2.5.2.2.1. Conventional synthesis. The Knoevenagel cyclocondensation of aliphatic aldehyde **56** with malononitrile **95** and **1** in ethanol at 20 °C in the presence of morpholine yielded the corresponding derivatives of 4-alkyl(cycloalkyl)-2-amino-3-cyano-4H-pyran **96** which are potential biologically active compounds. Their analogs are used in the treatment of cardiovascular diseases and CNS disorders, as well as for the protection




56,96	Ar	Yield %
a	4-Py	78
b	2-(5-CH ₃ C ₄ H ₂ O)	68
c	4-iPrO-C ₆ H ₄	90
d	3-CH ₃ O-4-iPrOC ₆ H ₃	74
e	3-CH ₃ O-4-(OCH ₂ CONH ₂)C ₆ H ₃	93

Scheme 49




Scheme 50



98,99,100	R	R ₁	Yield% (A/B/C)
a	Me	tert-Butyl	88/93/-
b	Me	Cyclohexyl	78/68/97
c	Et	Cyclohexyl	90/--/90

A : CH₂Cl₂ , r.t , 24 h
 B : Benzene, 80 °C, 5-6 h
 C : Acetone , r.t .

Scheme 51




56,101	Ar	Time/min (A/B/C)	Yield% (A/B/C)
a	-Ph	7/8/45	92/92/91
b	4-MeO-Ph	10/10/60	/89/85
c	4-NO ₂ -Ph	5/5/45	86/96/86
d	2,4-Cl-Ph	8/15/50	88/95/87
e	4-Cl-Ph	30/5/5/45	94/97/93
f	4-OH-Ph	12/30/60	91/90/85
g	4-Br-Ph	5/5/50	94/93/91
h	4-F-Ph	5/5/60	84/96/84
i	4-Me-Ph	8/8/60	88/90/85

Catalyst:

A : 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU).

B : 1-Butyl-3-methyl imidazolium hydroxide ([bmim]OH).

C : Tetrabutylammonium bromide (TBAB).

Scheme 52

56,102	Ar	Time/min (A/B/C/D)	Yield% (A/B/C/D)
a	-Ph	240/180/32/30	81/72/89/89
b	4-MeO-Ph	240/180/35/50	80/73/87/80
c	3-NO ₂ -Ph	240/180/30/45	93/88/91/80
d	4-NO ₂ -Ph	240/180/30/85	95/82/92/85
e	2,4-Cl-Ph	240/180/30/75	90/75/92/75
f	4-Cl-Ph	240/180/30/75	85/78/91/75


Catalyst:

A: Diammonium hydrogen phosphate (DAHP), r.t.


B: (S)-proline, reflux.

C: Magnesium oxide (MgO), reflux with stirring.


D: H₆P₂W₁₈O₆₂·18H₂O, reflux.**Scheme 53**



Scheme 54



Scheme 55



Scheme 56

of crops from herbicide damage (Dyachenko and Chernega, 2006; Dyachenko, 2005) (Scheme 47).


Klokot et al. (1999) described three component condensation of an aliphatic aldehyde **56** with malononitrile **95** and **1** at 20 °C in ethanol containing an equimolar amount of *N*-methylmorpholine led to the formation of the corresponding derivatives of 4-alkyl(cycloalkyl)-2-amino-3-cyano-4H-pyrans **96** (Scheme 48).

Shestopalov et al. (2005) noted that reaction of **1** with aromatic aldehyde **56** and malononitrile **95** in boiling ethanol in the presence of triethylamine as a catalyst gave substituted 2-amino-5-oxo-4,5-dihydropyrano[3,2-c]chromenes **96** in high yields (68–93%) (Scheme 49).

Okamoto et al. (1997, 1996) investigated that a reaction of **1**, pyruvaldehyde dimethyl acetal **97** and malononitrile **95** in boiling benzene containing piperidine gave 2-amino-3-cyano-4-dimethoxymethyl-4-methyl-5-oxo-4H,5H-pyrano[3,2-c]chromene **98** in 92% yield (Scheme 50).


The one-pot three-component chemoselective condensation reaction of substituted isocyanides **98** with stoichiometric amount of dialkyl acetylenedicarboxylates **99** in the presence of **1** proceeded spontaneously under different conditions led to a facile synthesis of highly functionalized corresponding dialkyl-2-(alkylamino)-5-oxo-4H,5H-pyrano-[4,3-b] pyran-3,4-dicarboxylates **100**, in moderate to good yields (68–97%) (Cravotto et al., 2011; Sarma et al., 2010; Tietze et al., 2001; Yavari et al., 2008) (Scheme 51).

3.2.5.2.2.2. *Using water.* The one-pot, three-component reaction of diversely substituted aromatic aldehydes **56** with malononitrile **95** and **1** in water under reflux in the presence of a catalyst (DBU, ([bmim]OH), TBAB, titanium dioxide, sodium tungstate or spinel zinc ferrite) gave, in all cases, the corresponding dihydropyrano[c]chromenes **102** in good to excellent yields (Das et al., 2014; Gong et al., 2009; Khodabakhshi and Baghernejad, 2014; Khodabakhshi et al., 2014; Khurana et al., 2010; Khurana and Kumar, 2009) (Scheme 52). It is worthwhile to note that the reaction is faster with aldehydes having electron-withdrawing groups (such as nitro group and halide) than that with electron-donating groups (such as methoxyl group and hydroxyl group).



56,105	a	b	c	d	e	f	h
R	H	2-Br	3-Me	3-OMe	3-Cl	4-Cl	4-Me
Yield %	94	92	85	83	78	75	81

Scheme 57



106,107	Ar	t/min	Yield%
a	4-CH ₃ OC ₆ H ₄	10	92
b	Piperonyl	7	89
c	2,5-(CH ₃ O) ₂ C ₆ H ₃	10	91
d	4-ClC ₆ H ₄	15	78

Scheme 58

Heravi et al. (2008) demonstrated an elegant protocol and eco-friendly route to the synthesis of 2-amino-4-aryl-5-oxo-4H,5H-pyrano[3,2-c]chromene-3-carbonitriles **101** via a three component reaction of aryl aldehydes **56**, malononitrile **95** and **1** using H₆P₂W₁₈O₆₂ · 18H₂O, which proceeds efficiently in aqueous ethanol under heating conditions. Many catalysts (DAHP or (S)-proline (Abdolmohammadi and Balalaie, 2007) or MgO (Seifi and Sheibani, 2008) or FeNi₃-SiO₂ (Nasseri and Sadeghzadeh, 2013)) have also been used as


another catalysts for this reaction (Scheme 53). Recently, Yao et al. (2013) found that 2-amino-4-aryl-5-oxo-4H,5H-pyrano[3,2-c]chromene-3-carbonitriles **101** could be prepared without catalyst in a mixed solvent of ethanol and water.

3.2.5.2.2.3. *Using ionic liquid.* 1,1,3,3-*N,N,N',N'*-Tetramethylguanidinium trifluoroacetate (TMGT) as an ionic liquid, efficiently promoted one-pot, three-component condensation of aldehydes **56**, alkynitriles **95** and **1** afforded pyran annulated heterocyclic systems **101** (Shaabani et al., 2005a) (Scheme 54).

3.2.5.2.2.4. *Microwave irradiation.* Kidwai and Saxena (2006) exploited an easier, practically convenient, novel, ecologically safe method for the synthesis of pyrano[3,2-c]benzopyran (R = phenyl, 4-chlorophenyl, 2-furyl, 3-indolyl, piperonyl, 2-chloro-3-quinolyl) **101** by three-component reaction of aldehyde **56**, malononitrile **95** and **1** in aqueous K₂CO₃ as a green catalyst under microwave heating. The observed yields and enhancement in reaction rates can be attributed to the uniform heating effect of microwave (Scheme 55).

3.2.6. 3,4-Dihydro-2H,5H-1-benzopyrano[4,3-b]pyran-5-one


Treatment of **1** with 1-aryl-2-[(dimethylamino)methyl]-2-propan-1-ones **102** in dimethylformamide in the presence of a catalytic amount of dimethylamine under reflux conditions afforded 3-benzoyl-3,4-dihydro-2H,5H-1-benzopyrano[4,3-b]pyran-5-ones **103** (Girreser et al., 1998) (Scheme 56).



R : Ph Me , 2-ClC₆H₄, 2-MeOC₆H₄ , 3-MeOC₆H₄ , 3-OHC₆H₄ , 3-NO₂C₆H₄ , 4-MeOC₆H₄ , 4-ClC₆H₄, 4-NO₂C₆H₄, 3,4-(Me)₂C₆H₃, 2-F-6-ClC₆H₃, 2,4-(Cl)₂C₆H₃ , 3-MeO-4-OHC₆H₃ , Furan-2-yl , Thiophene-2-yl , CH₃CH₂ , CH₃CH₂CH₂


R₁ : H, Me

Scheme 59



110,112	Ar	Yield %
a	Ph	83
b	4-Cl-C ₆ H ₄	74
c	4-NO ₂ -C ₆ H ₄	70
d	furyl	68

Scheme 60




n : 0,1,2

R₁ : iPr, iBu, Bn, MeR₂ : iPr, Et, Me, H


Cbz : benzyloxycarbonyl

Bn : Benzyl.

Scheme 61



Scheme 62





Scheme 63


Proline-promoted efficient enantioselective synthesis of 4-aryl-3,4-dihydro-2*H*,5*H*-pyrano[3,2-*c*]chromene-2,5-diones **105** from **1**, Meldrum's acid **104**, and benzaldehydes **56** (Yavari et al., 2008) (Scheme 57). The functionalized chromenes reported in this work may be considered as potentially

useful synthetic intermediates because they possess atoms with different oxidation states.


Khurana and Vij (2011) reported that the reaction of 5-arylidene Meldrum's acids **106** with **1** in the presence PEG-stabilized Ni nanoparticles underwent rapid and tandem



Scheme 64



Scheme 65



120,123	Ar	X	t /Sec. (A/B/C)	Yield% (A/B/C)
a	4-ClC ₆ H ₄	O	100/60/100	83/90/85
b	4-ClC ₆ H ₄	S	110/65/100	80/92/82
c	3-Furyl	O	100/75/85	82/95/84
d	3-Furyl	S	120/70/95	79/92/80
e	4-N(CH ₃) ₂ C ₆ H ₄	O	115/75/100	80/89/82
f	4-N(CH ₃) ₂ C ₆ H ₄	S	110/60/105	81/93/85
g	3-Indolyl	O	110/65/100	82/92/83
h	3-Indolyl	S	120/70/90	80/90/85
i	3-NO ₂ C ₆ H ₄	O	100/65/85	75/93/80
j	3-NO ₂ C ₆ H ₄	S	105/60/95	79/90/85

A : Silica gel
 B : Montmorillonite
 C : Acidic alumina.

Scheme 66

enol lactonization involving Michael addition to give 4-aryl-3,4-dihydropyrano[3,2-c]chromene-2,5-diones **107** (Scheme 58).

Three-component cyclocondensation of **1**, aldehydes **56**, and cyclic 1,3-dicarbonyl compounds **108** were prompted by ionic liquids 1,3-dimethyl-2-oxo-1,3-bis(4-sulfobutyl)imidazolidine-1,3-dium hydrogen sulfate ([DMDBSI].2HSO₄) in water to provide a novel series of 10,11-dihydro-chromeno[4,3-b]chromene-6,8(7*H*,9*H*)-dione derivatives **109** (Chen et al., 2011) (Scheme 59). Also, this reaction can be achieved using a catalytic amount of heteropolyacids or molecular iodine (Motamedi et al., 2012; Sun et al., 2012).

Cravotto et al. (2011) have disclosed a distinct improvement in the synthesis of pyranocoumarin derivatives through a pericyclic approach. This route offers a valid alternative to the standard industrial synthesis for 3,4-dihydropyrano[3,2-c]chromene-2,5-diones **112** were obtained by a hetero Diels–Alder (HDA) with inverse electron demand between **1**, aromatic aldehydes **110** (such as benzaldehyde, 4-chlorobenzaldehyde, 4-nitrobenzaldehyde, and 2-furancarboxaldehyde) and electron-rich alkenes **111** (Scheme 60).

Tietze et al. (2001) concluded that a multicomponent domino Knoevenagel/hetero-Diels–Alder reaction of **1** with an amino aldehyde **113** and benzyl enol ethers **114**, in toluene in the presence of catalytic amounts of ethylenediammonium diacetate (EDDA) and trimethyl orthoformate as dehydrating agent in an ultrasonic bath at 50 °C affords a benzyl-protected acetal **115** (Scheme 61).


Reaction of **1** with allene **116** in tetrahydrofuran in the presence of tetrakis(triphenylphosphine)palladium [Pd(PPh₃)₄]

afforded 3,4-dihydro-2,2-dimethyl-3-methylenepyrano[3,2-c]chromene-5(2*H*)-one **117** (Grigg et al., 2001) (Scheme 62).


3.2.7.4. 6*H*,11*H*-[2]Benzopyrano[4,3-*c*][1]benzopyran-11-one. The reaction of 2-(bromomethyl)phenyllead triacetate **118** with **1** carried out in the presence of a combination of *o*-phenanthroline-potassium *t*-butoxide (3:1) as a base for reductive coupling gave 6*H*,11*H*-[2]benzopyrano[4,3-*c*][1]benzopyran-11-one **119** (Ganina et al., 2005; Naumov et al., 2005) (Scheme 63).

3.2.8. Benzopyrano[4,3-*d*]pyrimidines

3.2.8.1. Using conventional heating. 4-Aryl-1,2,3,4-tetrahydro benzopyrano[4,3-*d*]pyrimidine-2,5-diones/pyrimidine-thione-5-ones **121** were prepared by reacting **1** with aromatic aldehydes **56** and urea/thiourea **120** in ethanol in the presence of concentrated hydrochloric acid, but this procedure requires prolonged heating of the reaction mixture and tedious workup



Scheme 68




56, 120, 124	X	Ar	t /min.	Yield %
a	O	Ph	2.6	94
b	S	Ph	22	92
c	O	3-indolyl	2.9	90
e	S	3-indolyl	3.1	88
f	O	1,3-2H-benzo dioxol-5-yl	3.3	93
g	S	1,3-2H-benzo dioxol-5-yl	3.4	95
h	O	2-chloroquin-olin-3-yl	4.1	86
i	S	2-chloroquin-olin-3-yl	4.3	87

Scheme 67


and gives low yields of the products (Brahmbhatt et al., 1999) (Scheme 64).

3.2.8.2. *Solution phase synthesis.* Due to short reaction times, high yields and easy work-up procedures combined with the use of the multiple synthesizer, Sabitha et al. (2003) found that



127,128	Ar	Yield %
a	Ph	75
b	2-FC ₆ H ₄	67
c	4-NO ₂ C ₆ H ₄	72

Scheme 69




129,130	Ar	Yield %
a	4-MeC ₆ H ₄	75
b	4-MeOC ₆ H ₄	71

Scheme 70

three component condensation of **1**, an aromatic aldehyde **56**, and urea (thiourea) **120** in refluxing acetonitrile containing a catalytic amount of VCl₃ gave the dihydropyrimidinones **122** with high purity > 95%. (Scheme 65).


3.2.8.3. *Microwave irradiation.* Acidic solid support coupled with microwave is an appropriate method for the synthesis of 4-substituted benzopyrano[4,3-d]pyrimidine derivatives **123** in a few seconds with improved yield as compared to conventional heating. The main advantage being that solid supports do not absorb microwaves at 2450 MHz, so are not an obstacle for the transition of microwaves to the reactants. In addition, the limitations of the MWI assisted reaction in solvents, namely, the development of high pressure and the need for specialized sealed vessels are circumvented via this solid state technique which enables organic reactions to occur rapidly at atmospheric pressure. Montmorillonite K10 clay is a better support as compared to silica gel and acidic alumina in terms of yield and time for the synthesis of benzopyranopyrimidines **123** (Kidwai and Sapra, 2002) (Scheme 66).

Kidwai et al. (2006) developed a highly efficient environmentally benign method is proposed for the synthesis of benzopyranopyrimidines **124** apart from acidic solid support by reaction of **1** with aldehydes **56** and urea or thiourea **120** in the absence of a solvent under microwave irradiation in a few minutes gave the desired products in 90–95% yields. The



134,135	a	b
R		
Yield %	24	11

Scheme 72



131,133	a	b	c	d	e	f
Ar	Ph	4-Cl-Ph	4-Br-Ph	4-CH ₃ -Ph	3-NO ₂ -Ph	4-NO ₂ -Ph
Yield %	91	93	94	90	90	91

Scheme 71

proposed procedure utilizes neither solvents nor solid support nor acid catalyst, requires no special equipment, improves the product yield and shortens the reaction time and minimizes hazardous pollution (Scheme 67).

3.2.9. Benzopyranobenzothiazinones

One of the most successful strategies for constructing of 6,12-dihydro-1-benzopyrano[3,4-b][1,4]benzothiazin-6-one **126**, new agents with estrogenic activity mediated by estrogen receptors (ER), is the condensation and oxidative cyclization of substituted 2-aminobenzenethiol **125** with **1** in dimethyl sulfoxide at 140–150 °C (Aguirre-Pranzoni et al., 2011; Anary-Abbasinejad et al., 2008; Gupta et al., 2004; Gupta and Gupta, 2009) (Scheme 68).

3.2.10. Chromeno[3,4-e][1,3]oxazine

Vovk et al. (2007) have described the reaction of **1** with 1-chlorobenzyl isocyanates **127** occurred in anhydrous toluene at 60 °C, did not require the presence of an organic base, and led to the formation of 4-aryl-3,4-dihydro-2*H*,5*H*-chromeno[3,4-e][1,3]oxazine-2,5-diones **128** in 67–75% yields (Scheme 69).

However, 1-aryl-2,2,2-trifluoro-1-chloroethyl isocyanates **129** reacted with **1** in the presence of triethylamine in toluene gave 2-aryl-2-trifluoromethyl-2,3-dihydro-4*H*,5*H*-chromeno[3,4-e][1,3]oxazine-4,5-diones **130** (Vovk et al., 2007) (Scheme 70).

Hassanabadi et al. (2011) concluded that the reaction of acid chlorides **131** and ammonium thiocyanate **132** with **1** in the presence of *N*-methylimidazole led to oxazine derivatives **133** in excellent yields. The present procedure has the advantage that the reaction is performed under neutral conditions

and the starting material can be used without any activation or modification (Scheme 71).

3.2.11. Pyrano[3,4-e][1,4,3]oxathiazines

Cablewski et al. (2007) investigated the regioselective reaction of **1** with *N,N*-dialkyl-*N'*-chlorosulfonylchloro formamides **134** in the presence of *N,N'*-dimethyl-*N,N'*-propylene urea (DMPU) afforded 3-dialkylamino-1,1,8-trioxo-1*H*-6-pyrano[3,4-e][1,4,3]oxathiazines **135** (Scheme 72).


3.3. [7-6] Ring system

3.3.1. Benzopyrano[1,3]diazepines

Kidwai et al. (2007) have shown that a simple and facile synthesis of 7-arylbenzopyrano[1,3]diazepines **137** was accomplished by the treatment of **1**, aromatic/heteroaromatic aldehydes **56**, and cyanoguanidine **136** using molecular iodine as a novel catalyst in acetonitrile under reflux. This method not only provides an excellent complement to benzopyrano[1,3]diazepines **137** but also avoids the multistep and harsh reaction conditions (Scheme 73).

4. Conclusion

The present review is a summary of progress in 4-hydroxycoumarin as a versatile synthetic building block to the synthesis of all sorts of heterocycles or fused heterocycles. Even though a lot of progress has reported, We hope more astonishing applications of this compound will be revealed in the near future, and that this review may light a candle in organic chemistry.



Scheme 73

References

- Abd El-Aziz, A.S., Mohamed, H.M., Mohammed, S., Zahid, S., Ata, A., Bedair, A.H., El-Agrody, A.M., Harvey, P.D., 2007. Synthesis of novel coumarin and benzocoumarin derivatives and their biological and photophysical studies. *J. Heterocycl. Chem.* 44 (6), 1287–1301.
- Abd El-Wahab, A.H.F., Al-Fifi, Z.I.A., Bedair, A.H., Ali, F.M., Halawa, A.H.A., El-Agrody, A.M., 2011. Synthesis, reactions and biological evaluation of some new naphtho[2,1-b]furan derivatives bearing a pyrazole nucleus. *Molecules* 16 (1), 307–318.
- Abdolmohammadi, S., 2014. Silica supported $Zr(HSO_4)_4$ catalyzed solvent-free synthesis of [1]benzopyrano[4,3-b][1]benzopyran-6-ones and xanthenones. *J. Org. Chem.* 11 (5), 350–355.
- Abdolmohammadi, S., Balalaie, S., 2007. Novel and efficient catalysts for the one-pot synthesis of 3,4-dihydropyrano[c]chromene derivatives in aqueous media. *Tetrahedron Lett.* 48 (18), 3299–3303.
- Abdou, M.M., 2017. Utility of 4-hydroxythiocoumarin in organic synthesis. *Arab. J. Chem.* 10, S3955–S3961.
- Abdou, M.M., 2013b. New Azo Disperse Dyes Derived from 3-(Hydroxyphenyl)-2-Pyrazolin-5-One. German publisher LAP (Lambert Academic Publishing).
- Abdou, M.M., 2013c. Azo Disperse Dyes with 2-Pyrazolin-5-Ones for Dyeing Polyester Fabrics. German publisher LAP (Lambert Academic Publishing).
- Abdou, M.M., 2017. Chemistry of 4-hydroxy-2(1H)-quinolone. Part 1: Synthesis and reactions. *Arab. J. Chem.* 10, S3324–S3337.
- Abdou, M.M., 2018. Chemistry of 4-hydroxy-2(1H)-quinolone. Part 2 as synthons in heterocyclic synthesis. *Arab. J. Chem.* 10, 1061–1071.
- Abdou, M.M., 2019. Recent advances in 4-hydroxycoumarin chemistry. Part 1: Synthesis and reactions. *Arab. J. Chem.* 12, 88–121.
- Abdou, M.M., 2017a. Chemistry of 3-acetyl-4-hydroxycoumarin. *Arab. J. Chem.* 10 (2), S3664–S3675.
- Abdou, M.M., 2017b. 3-Acetyl-4-hydroxycoumarin: Synthesis, reactions and applications. *Arab. J. Chem.* 10 (2), S3664–S3675.
- Abdou, M.M., Bondock, S., El-Desouky, E.I., Metwally, M.A., 2012. Synthesis, structure elucidation and application of some new azo disperse dyes derived from 4-hydroxycoumarin for dyeing polyester fabrics. *Am. J. Chem.* 2 (6), 347–354.
- Abdou, M.M., Bondock, S., El-Desouky, S.I., Metwally, M.A., 2013. Synthesis, spectroscopic studies and technical evaluation of novel disazo disperse dyes derived from 3-(2-hydroxyphenyl)-2-pyrazolin-5-ones for dyeing polyester fabrics. *Am. J. Chem.* 3 (3), 59–67.
- Adib, M., Mahdavi, M., Bagherzadeh, S., Bijanzadeh, H.R., 2009. An efficient and direct solvent-free synthesis of naphtho[1,2-b]furans, naphtho[2,1-b]furans, and furo[3,2-c]chromenes. *Synlett* 15, 2542–2544.
- Aggarwal, K., Khurana, J.M., Saluja, P., 2013. One-pot synthesis of biologically important spiro-2-amino-4H-pyrans, spiroacenaphthylenes, and spirooxindoles using DBU as a green and recyclable catalyst in aqueous medium. *Synth. Commun.* 43 (24), 3239–3246.
- Aguirre-Pranzoni, C.B., Furque, G.I., Ardanaz, C.E., Pacciaroni, A., Sosa, V., Tonn, C.E., Kurina-Sanz, M., 2011. Biotransformation of dihydrocoumarin by *Aspergillus niger* ATCC 11394. *Arkivoc* 7, 170–181.
- Ahadi, S., Zolghadr, M., Khavasi, H.R., Bazgir, A., 2013. A diastereoselective synthesis of pyrano fused coumarins via organocatalytic three-component reaction. *Org. Biomol. Chem.* 11 (2), 279–286.
- Ahadi, S., Zolghadr, M., Khavasi, H.R., Bazgir, A., 2014. An organocatalytic three-component synthesis of chromeno [4, 3-b] chromenones. *J. Iran. Chem. Soc.* 11 (1), 155–162.
- Altieri, E., Cordaro, M., Grassi, G., Risitano, F., Scala, A., 2010. Regio and diastereoselective synthesis of functionalized 2,3-dihydrofuro[3,2-c]coumarins via a one-pot three-component reaction. *Tetrahedron* 66 (49), 9493–9496.
- Anary-Abbasinejad, M., Anaraki-Ardakani, H., Hassanabadi, A., 2008. P_2O_5 -Hexamethyl disiloxane (HMDS): an efficient system to induce the three-component reaction of enolic systems, aromatic aldehydes, and acetonitrile. *Synth. Commun.* 38 (21), 3706–3716.
- Angeleska, S., Kefalas, P., Detsi, A., 2013. Crude peroxidase from onion solid waste as a tool for organic synthesis. Part III: Synthesis of tetracyclic heterocycles (coumestans and benzofuroquinolones). *Tetrahedron Lett.* 54 (19), 2325–2328.
- Appendino, G., Cravotto, G., Palmisano, G., Annunziata, R., 1998. Chemistry of coumarin derivatives. IX. Synthesis of fercoprolone, a degraded prenylated coumarin. *Tetrahedron* 54 (36), 10819–10826.
- Azizi, N., Dezfooli, S., Mahmoudi, H.M., 2014. Greener synthesis of spirooxindole in deep eutectic solvent. *J. Mol. Liq.* 194, 62–67.
- Baichurin, R.I., Baichurina, L.V., Aboskalova, N.I., Berestovitskaya, V.M., 2013. Synthesis of benzoyl- and ethoxycarbonyl-containing dihydrofurocoumarins. *Russ. J. Gen. Chem.* 83 (9), 1787–1789.
- Bakthadoss, M., Sivakumar, G., 2014. Highly stereo and chemoselective synthesis of tetra and pentacyclic frameworks using Solid-State Melt Reaction (SSMR). *Tetrahedron Lett.* 55 (10), 1765–1770.
- Banard, D.L., Xu, Z.Q., Stowell, V.D., Yuan, H.W., Smee, D.F., Samy, R., Ridwell, R.W., Nielson, M.K., Sun, L.H., Cao, H., Li, A.L., Quint, C., Deignan, J., Crabb, J., Flavin, M.T., 2002. Coumarins and pyranocoumarins: potential novel pharmacophores for inhibition of measles virus replication. *Antiviral Chem. Chemother.* 13, 39–59.
- Bandyopadhyay, C.A., Maiti, S., Panja, S.K., 2010. One-pot synthesis of the 1-benzopyrano [2,3-b]pyridine moiety from 2-(alkyl/aryl amino)-4-oxo-4H-1-benzopyran-3-carbaldehyde. *J. Heterocycl. Chem.* 47 (4), 973–981.
- Bar, G., Bini, F., Parsons, A.F., 2003. CAN-mediated oxidative free radical reactions in an ionic liquid. *Synth. Commun.* 33 (2), 213–222.
- Bedair, A.H., El-Hady, N.A., El-Latif, M.S. Abd., Fakery, A.H., El-Agrody, A.M., 2000. 4-Hydroxycoumarin in heterocyclic synthesis Part III. Synthesis of some new pyrano[2,3-d]pyrimidine, 2-substituted [1,2,4]triazolo[1,5-c]pyrimidine and pyrimido [1,6-b][1,2,4] triazine derivatives. *Farmaco* 55 (11–12), 708–714.
- Bondock, S., Khalifa, W., Fadda, A.A., 2011. Synthesis and antimicrobial activity of some new 4-hetarylpyrazole and furo [2,3-c]pyrazole derivatives. *Eur. J. Med. Chem.* 46 (6), 2555–2561.
- Brahmbhatt, D.I., Raolji, G.B., Pandya, S.U., Pandya, U.R., 1999. Synthesis of some 4-aryl-1,2,3,4-tetrahydrobenzopyrano[4,3-d]pyrimidine-2,5-diones/pyrimidine-2-thione-5-ones. *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.* 38B (7), 839–842.
- Cablewski, T., Francis, C.L., Liepa, A.J., 2007. N,N-Dialkyl-N'-chlorosulfonyl chloro formamidines in heterocyclic synthesis. IV. 3-Dialkylamino-1,1,8-trioxo-1H-12,6-pyrano[3,4-e][1,4,3]oxathiazines. *Aust. J. Chem.* 60 (2), 113–119.
- Cadierno, V., Diez, J., Gimeno, J., Nebra, N., 2008. Ruthenium/TFA-catalyzed coupling of activated secondary propargylic alcohols with cyclic 1,3-diones: furan versus pyran ring formation. *J. Org. Chem.* 73 (15), 5852–5858.
- Chen, Z., Zhu, Q., Su, W., 2011. A novel sulfonic acid functionalized ionic liquid catalyzed multicomponent synthesis of 10, 11-dihydrochromeno[4,3-b]chromene-6,8(7H,9H)-dione derivatives in water. *Tetrahedron Lett.* 52, 2601–2604.
- Chen, Zhi-Wei., Su, Wei-Ke., Wang, Zhi-Hua., Zhang, Na., 2013. An efficient synthesis of novel chromeno[3',4':5,6]pyrano[2,3-b] indole derivatives. *Chin. Chem. Lett.* 24 (3), 199–201.
- Cravotto, G., Nano, G.M., Palmisano, G., Pilati, T., Tagliapietra, S., 2011. A library of pyranocoumarin derivatives via a one-pot three-component hetero Diels-Alder reaction. *J. Heterocycl. Chem.* 38 (4), 965–971.
- Darehkordi, A., Karimi-Taleghani, Z., Pouralimardan, O.A., Peng, S., Gao, T., Sun, S., Peng, Y., Wu, M., Guo, H., Wang, J., 2014.

- Palladium-catalyzed [2 + 2 + 1] oxidative annulation of 4-hydroxycoumarins with unactivated internal alkynes: Access to spiro cyclopentadiene-chroman-2,4-dione complexes. *Adv. Synth. Catal.* 356 (2–3), 319–324.
- Das, P., Dutta, A., Bhaumik, A., Mukhopadhyay, C., 2014. Heterogeneous ditopic ZnFe_2O_4 catalyzed synthesis of 4 H-pyrans: further conversion to 1,4-DHPs and report of functional group interconversion from amide to ester. *Green Chem.* 16, 1426–1435.
- Davarpanah, J., Kiasat, A.R., 2014. Covalently anchored n-propyl-4-aza-1-azoniabicyclo[2.2.2]octane chloride on SBA-15 as a basic nanocatalyst for the synthesis of pyran heterocyclic compounds. *RSC Adv.* 4 (9), 4403–4412.
- Dekamin, M.G., Eslami, M., Maleki, A., 2013. Potassium phthalimide-N-oxyl: a novel, efficient, and simple organocatalyst for the one-pot three-component synthesis of various 2-amino-4H-chromene derivatives in water. *Tetrahedron* 69 (3), 1074–1085.
- Dyachenko, V.D., 2005. Isovaleric aldehyde in the synthesis of 4-isobutyl-substituted pyridine-2(1H)-thiones, 4H-pyrans, and 1,3-cyclohexadiene. *Russ. J. Gen. Chem.* 75 (3), 440–446.
- Dyachenko, V.D., Chernega, A.N., 2006. Aliphatic aldehydes in multicomponent syntheses of 4-alkyl-substituted partially hydrogenated quinolines, fused 4H-pyrans, and 2-amino-4-ethyl-5-methylbenzene-1,3-dicarbonitrile. *Russ. J. Org. Chem.* 42 (4), 567–576.
- Dyachenko, V.D., Rusanov, E.B., 2006. Isopropylidenemalononitrile in the synthesis of 2-amino-4,6,6-trimethylcyclohexa-2,4-diene-1,1,3-tricarbonitrile, 6-amino-2-(4-methoxybenzoylmethylsulfanyl)-4,4-dimethyl-1,4-dihydropyridine-3,5-dicarbonitrile, and fused 2-aminopyrans via Michael reactions. *Russ. J. Org. Chem.* 42 (9), 1374–1379.
- El-Agrody, A.M., Abd El-Latif, M.S., Fakery, A.H., Bedair, A.H., 2000. Heteroaromatization with 4-hydroxycoumarin Part I: Synthesis of some new pyranocoumarins and coumarinopyranopyrimidines. *J. Chem. Res. (S)* 1, 26–27.
- Eskandari, K., Karami, B., Khodabakhshi, S., 2013. A new application of Mohr's salt as a cheap and powerful catalyst for synthesis of novel pyranocoumarins. *Lett. Org. Chem.* 10 (2), 105–110.
- Froggett, J.A., Hockley, M.H., Titman, R.B., 1997. Novel preparation of 1-aryl-3-(2-hydroxyphenyl)-2-pyrazolin-5-ones and their conversion into 2-aryl-4-methyl[1] benzopyrano[4,3-c]pyrazol-3(2H)-ones. *J. Chem. Res. (S)* 1, 30–31.
- Ganina, O.G., Zamotaeva, S.G., Nosarev, M.A., Kosenkova, O.V., Naumov, M.I., Shavrin, A.S., Finet, J.P., Fedorov, A.Y., 2005. 2-(Azidomethyl)phenylboronic acid in the synthesis of isoquinoline derivatives. *Russ. Chem. Bull.* 54 (7), 1606–1611.
- Garella, D., Barge, A., Upadhyaya, D., Rodriguez, Z., Cravotto, G., Palmisano, G., 2010. Fast, solvent-free, microwave-promoted Friedländer annulation with a reusable solid catalyst. *Synth. Commun.* 40 (1), 120–128.
- Ghandi, M., Taghi Nazeri, M., Kubicki, M., 2013. An efficient one-pot, regio- and stereoselective synthesis of novel pentacyclic-fused pyrano [3, 2, c] chromenone or quinolinone benzosultone derivatives in water. *Tetrahedron* 69 (24), 4979–4989.
- Ghosh, A., Khan, A.T., 2014. Synthesis of dihydrochromeno[4,3-b]pyrazolo[4,3-c]pyridin-6(7H)-ones involving one-pot three-component tandem Knoevenagel-Michael reaction catalyzed by n-tetrabutylammonium tribromide (TBATB). *Tetrahedron Lett.* 55 (12), 2006–2009.
- Girreser, U., Heber, D., Schutt, M., 1998. 3-Benzoyl-3,4-dihydro-2H,5H-1-benzopyrano[4,3-b]pyran-5-ones by condensation of 4-hydroxycoumarins with enone Mannich bases. *J. Heterocycl. Chem.* 35 (6), 1455–1460.
- Golabi, S.M., Nematollahi, D., 1997a. Electrochemical study of 3,4-dihydroxybenzoic acid and 4-tert-butylcatechol in the presence of 4-hydroxycoumarin. Application to the electro-organic synthesis of coumestan derivatives. *J. Electroanal. Chem.* 430 (1–2), 141–146.
- Golabi, S.M., Nematollahi, D., 1997b. Electrochemical study of catechol and some 3-substituted catechols in the presence of 4-hydroxycoumarin: application to the electro-organic synthesis of new coumestan derivatives. *J. Electroanal. Chem.* 420 (1–2), 127–134.
- Golzar, N., Hasaninejad, A., Zare, A., 2013. One-pot, four-component synthesis of novel spiro[indeno[2,1-b]quinoxaline-11,4'-pyran]-2'-amines. *J. Heterocycl. Chem.* 50 (3), 608–614.
- Gong, K., Wang, H.L., Luo, J., Liu, Z.L., 2009. One-pot synthesis of polyfunctionalized pyrans catalyzed by basic ionic liquid in aqueous media. *J. Heterocycl. Chem.* 46 (6), 1145–1150.
- Grigg, R., Kongathip, N., Kongathip, B., Luangkamin, S., Dondas, H.A., 2001. Palladium catalysed reactions of allene with active methylene pronucleophiles. C-1, 3-dienylmethyl derivatives and their Diels-Alder reactions. *Tetrahedron* 57 (44), 9187–9197.
- Gu, Y., Barrault, J., Jerome, F., 2009. Trapping of active methylene intermediates with alkenes, indoles or thiols: towards highly selective multicomponent reactions. *Adv. Synth. Catal.* 351 (18), 3269–3278.
- Gupta, V., Gupta, A., 2009. An efficient synthesis of novel 6,12-dihydro(1)benzopyrano[3,4-b][1,4] benzothiazin-6-one. *Heterocycl. Commun.* 15 (3), 225–227.
- Gupta, R., Shrivastava, V., Gupta, A., Verma, P.S., Gupta, R.R., 2004. One-pot syntheses of heterocycles containing 1,4-thiazine nucleus. *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.* 43B (2), 413–416.
- Hamdi, N., Lidrissi, C., Saoud, M., Nieves, A., Romerosa, Zarrouk, H., 2006. Synthesis of some new biologically active coumarin derivatives. *Chem. Heterocycl. Compd.* 42 (3), 320–325.
- Hassanabadi, A., Mosslemin, M.H., Tadayonfar, S.E., 2011. Three-component reaction between 4-hydroxycoumarin, ammonium thiocyanate and acid chlorides in the presence of N-methylimidazole under solvent-free conditions. *J. Chem. Res.* 35 (1), 29–31.
- Heravi, M.M., Jani, B.A., Derikvand, F., Bamoharram, F.F., Oskooie, H.A., 2008. Three component, one-pot synthesis of dihydropyran[3,2-c]chromene derivatives in the presence of $\text{H}_2\text{P}_2\text{W}_{18}\text{O}_{62}\cdot 18\text{H}_2\text{O}$ as a green and recyclable catalyst. *Catal. Commun.* 10 (3), 272–275.
- Hossain, E., Hussain, S., Molla, A., 2013. Multicomponent domino reactions: borax catalyzed synthesis of highly functionalised pyran-annulated heterocycles. *RSC Adv.* 3 (44), 21517–21523.
- Hossaini, Z., Charati, F.R., Ghasemian, M., 2013. Synthesis of functionalized phosphonates and chromenes via catalyst-free multicomponent reactions in water. *Phosphorus, Sulfur Silicon Relat. Elem.* 188 (5), 555–560.
- Huang, C.N., Kuo, P.Y., Lin, C.H., Yang, D.Y., 2007. Synthesis and characterization of 2H-pyrano[3,2-c]coumarin derivatives and their photochromic and redox properties. *Tetrahedron* 63 (40), 10025–10033.
- Itoh, K., Kanemasa, S., 2003. A new method for enol lactone synthesis by a Michael addition/cyclization sequence. *Tetrahedron Lett.* 44 (9), 1799–1802.
- Itoh, K., Hasegawa, M., Tanaka, J., Kanemasa, S., 2005. Enantioselective enol lactone synthesis under double catalytic conditions. *Org. Lett.* 7 (6), 979–981.
- Jaggavarapu, S.R., Kamalakaran, A.S., Jalli, V.P., Gangisetty, S.K., Ganesh, M.R., Gaddamanugu, G., 2014. Facile eco-friendly synthesis of novel chromeno [4, 3-b] pyridine-2, 5-diones and evaluation of their antimicrobial and antioxidant properties. *J. Chem. Sci.* 126 (1), 187–195.
- Jakse, R., Svete, J., Stanovnik, B., Golobic, A., 2004. Application of alkyl 3-dimethylamino-2-(1H-indol-3-yl)propenoates in the synthesis of 3-heteroarylindoles. *Tetrahedron* 60 (21), 4601–4608.
- Jashari, A., Imeri, F., Ballazhi, L., Shabani, A., Mikhova, B., Draeger, G., Popovski, E., Huwiler, A., 2014. Synthesis and cellular characterization of novel isoxazolo- and

- thiazolohydrazinylidene-chroman-2,4-diones on cancer and non-cancer cell growth and death. *Bioorg. Med. Chem.* 22 (9), 2655–2661.
- Jeon, M.K., Kim, K., 1999. Synthesis of new 5-alkylidene-4-chloro-5H-1,2,3-dithiazoles and their stereochemistry. *Tetrahedron* 55 (32), 9651–9667.
- Jin, S.S., Wang, H., Guo, H.Y., 2013. Ionic liquid catalyzed one-pot synthesis of novel spiro-2-amino-3-phenylsulfonyl-4H-pyran derivatives. *Tetrahedron Lett.* 54 (19), 2353–2356.
- Jukic, L., Svete, J., Stanovnik, B., 2001a. Reactions of alkyl (Z)-2-[(E)-2-ethoxycarbonyl-2-(2-pyridinyl)-ethenyl]amino-3-dimethylaminopropenoates with C- and N-nucleophiles. The synthesis of fused pyranones, pyridinones and pyrimidinones. *J. Heterocycl. Chem.* 38 (4), 869–876.
- Jukic, L., Svete, J., Stanovnik, B.S., 2001b. Reactions of alkyl (Z)-2-[(E)-2-cyano-2-(2-pyridinyl) ethenyl]amino-3-dimethylaminopropenoates with C- and N-nucleophiles. The synthesis of fused 2H,5H-pyran-2,5-diones, 4H-pyrimidin-4-ones, and 1-heteroaryl-1H-imidazole-4-carboxylates. *Heterocycles* 55 (4), 705–727.
- Jung, J.C., Park, O.S., 2009. Synthetic approaches and biological activities of 4-hydroxycoumarin derivatives. *Molecules* 14 (11), 4790–4803.
- Karami, B., Khodabakhshi, S., Eskandari, K., 2013. Regiospecific synthesis of novel furo [4, 5-c] coumarins in a one-pot reaction. *Synlett* 24 (08), 998–1000.
- Khan, A.T., Lal, M., Ali, S., Khan, M.M., 2011. One-pot three-component reaction for the synthesis of pyran annulated heterocyclic compounds using DMAP as a catalyst. *Tetrahedron Lett.* 52 (41), 5327–5332.
- Khan, A.T., Lal, M., Basha, R.S., 2013. Regio- and diastereoselective synthesis of trans-2, 3-dihydrofuran derivatives in an aqueous medium. *Synthesis* 45 (03), 406–412.
- Khan, M.N., Pal, S., Karamthulla, S., Choudhury, L.H., 2014. Imidazole as organocatalyst for multicomponent reactions: diversity oriented synthesis of functionalized hetero- and carbocycles using in situ-generated benzylidenemalononitrile derivatives. *RSC Adv.* 4 (8), 3732–3741.
- Khodabakhshi, S., Baghernejad, M., 2014. Sodium tungstate: the first application as non-oxidative catalyst in organic synthesis. *J. Chin. Chem. Soc.* 61 (5), 521–524.
- Khodabakhshi, S., Karami, B., Eskandari, K., Hoseini, S.J., 2014. Titanium dioxide nanowires as green and heterogeneous catalysts for the synthesis of novel pyranocoumarins. *Comptes Rendus Chim.* 17 (1), 35–40.
- Khurana, J.M., Kumar, S., 2009. Tetrabutylammonium bromide (TBAB): a neutral and efficient catalyst for the synthesis of biscoumarin and 3,4-dihydropyranoc[chromene] derivatives in water and solvent-free conditions. *Tetrahedron Lett.* 50 (28), 4125–4127.
- Khurana, J.M., Vij, K., 2011. Nickel nanoparticles catalyzed chemoselective Knoevenagel condensation of Meldrum's acid and tandem enol lactonizations via cascade cyclization sequence. *Tetrahedron Lett.* 52 (28), 3666–3669.
- Khurana, J.M., Nand, B., Saluja, P., 2010. DBU: a highly efficient catalyst for one-pot synthesis of substituted 3,4-dihydropyranoc[3,2-c]chromenes, dihydropyranoc[4,3-b]pyranes, 2-amino-4H-benzo[h]chromenes and 2-amino-4H benzo[g]chromenes in aqueous medium. *Tetrahedron* 66 (30), 5637–5641.
- Kidwai, M., Sapra, P., 2002. An efficient synthesis of benzopyranopyrimidines using inorganic solid support. *Synth. Commun.* 32 (11), 1639–1645.
- Kidwai, M., Saxena, S., 2006. Convenient preparation of pyranobenzopyranes in aqueous media. *Synth. Commun.* 36 (18), 2737–2742.
- Kidwai, M., Saxena, S., Mohan, R.A., 2005. New method for the synthesis of heterocycles from o-phenylenediamine. *J. Korean Chem. Soc.* 49 (3), 288–291.
- Kidwai, M., Saxena, S., Mohan, R., 2006. Environmentally benign synthesis of benzo pyranopyrimidines. *Russ. J. Org. Chem.* 42 (1), 52–55.
- Kidwai, M., Bansal, V., Mothra, P., 2007. Molecular iodine: a highly efficient catalyst for the synthesis of 7-arylbenzopyranoc[1,3]diazepines in non-protic solvents. *J. Mol. Catal. A: Chem.* 266 (1–2), 43–46.
- Kislyi, V.P., Nesterov, V.N., Shestopalov, A.M., Semenov, V.V., 1999. Heterocycles with a β -nitroenamine fragment. 2. Synthesis of 2-amino-3-nitropyranoc[3,2-c]pyrans and 2-amino-3-nitropyranoc[3,2-c]chromenes from nitroacetonitrile. *Russ. Chem. Bull.* 48 (6), 1135–1138.
- Klokol, G.V., Krivokolysko, S.G., Dyachenko, V.D., Litvinov, V.P., 1999. Aliphatic aldehydes in the synthesis of condensed 4-alkyl(-cycloalkyl)-2-amino-3-cyano-4H-pyrans. *Chem. Heterocycl. Compd.* 35 (10), 1183–1186.
- Kobayashi, K., Sakashita, K., Akamatsu, H., Tanaka, K., Uchida, M., Uneda, T., Kitamura, T., Morikawa, O., Konishi, H., 1999. CAN-mediated formation of furopyranones and furoquinolinone. *Heterocycles* 51 (12), 2881–2892.
- Kolos, N.N., Gosalishvili, L.L., Sivokon, E.N., Knyazeva, I.V., 2009. Reaction of 4-hydroxycoumarin with arylglyoxals and ureas. *Russ. J. Org. Chem.* 45 (1), 119–125.
- Kovregin, A.N., Sizov, A.Y., Ermolov, A.F., 2001. Reactions of 1,3,3,3-tetrafluoro-2-methoxycarbonyl propenylsulfenyl chloride with aromatic and heterocyclic compounds. *Russ. Chem. Bull.* 50 (7), 1255–1258.
- Lamani, R.S., Shetty, N.S., Kamble, R.R., Khazi, I.A.M., 2009. Synthesis and antimicrobial studies of novel methylene bridged benzisoxazolyl imidazo[2,1-b][1,3,4]thiadiazole derivatives. *Eur. J. Med. Chem.* 44 (7), 2828–2833.
- Lee, Y.R., Kim, B.S., 1997. A facile synthesis of dihydrofurans utilizing silver(I)/celite promoted oxidative cycloaddition of 1,3-dicarbonyl compounds to alkenes. *Tetrahedron Lett.* 38 (12), 2095–2098.
- Lee, Y.R., Byun, M.W., Kim, B.S., 1998. Efficient one-step synthesis of 2-arylfurans by ceric ammonium nitrate (CAN)-mediated cycloaddition of 1,3-dicarbonyl compounds to alkynes. *Bull. Korean Chem. Soc.* 19 (10), 1080–1083.
- Lee, Y.R., Kim, B.S., Kim, D.H., 2000. Ceric ammonium nitrate (CAN)-mediated oxidative cycloaddition of 1,3-dicarbonyls to conjugated compounds. Efficient synthesis of dihydrofurans, dihydro furocoumarins, dihydrofuroquinolinones, dihydrofuro phenalenones, and furonaphtho quinone natural products. *Tetrahedron* 56 (45), 8845–8853.
- Lee, C.J., Jang, Y.J., Wu, Z.Z., Lin, W., 2012. Preparation of functional phosphorus zwitterions from activated alkanes, aldehydes, and tributylphosphine: synthesis of polysubstituted furo [3, 2-c] coumarins. *Org. Lett.* 14 (7), 1906–1909.
- Leutbecher, H., Conrad, J., Klaiber, I., Beifuss, U., 2005. O-Heterocycles via laccase-catalyzed domino reactions with O₂ as the oxidant. *Synlett* 20, 3126–3130.
- Leutbecher, H., Greiner, G., Amann, R., Beifuss, U., Conrad, J., Stolz, A., 2011. Laccase-catalyzed phenol oxidation. Rapid assignment of ring-proton deficient polycyclic benzofuran regioisomers by experimental ¹H–¹³C long-range coupling constants and DFT-predicted product formation. *Org. Biomol. Chem.* 9 (8), 2667–2673.
- Magiatis, P., Melliou, E., Skaltsounis, A.L., Mitaku, S., Leonce, S., Reard, P., Pierre, A., Atassi, G., 1998. Synthesis and cytotoxic activity of pyranocoumarins of the seselin and xanthyletin series. *J. Nat. Prod.* 61 (8), 982–986.
- Mahdavinia, G.H., Peikarporsan, S., 2013. Cyclization of α,α' -bis(substituted-benzylidene) cyclohexanones and 4-hydroxycoumarin: synthesis of 11-benzylidene-8,9,10,11-tetrahydro-7-phenyl-6H,7H-chromeno[4,3-b]chromen-6-ones as new pyranochromene derivatives. *Monatsh. Chem.* 144 (3), 415–419.

- Mahmoud, M.R., El-Bassiouny, F.A., Azab, M.E., El-Kady, M.Y., Rashed, H.M., 2009. Heteroannulation of chromene derivatives. Synthesis of chromeno[4,3-e]indazolone, chromeno[4,3-f]quinazoline and pyrano[3,2-c]chromene derivatives. *J. Chem. Res.* 1, 41–45.
- Mail, R.S., Joshi, P.P., Sandhu, P.K., Manekar-Tilve, A.J., 2002. Efficient syntheses of 6-prenylcoumarins and linear pyranocoumarins: total synthesis of suberosin, toddaculin, O-methylapigravin (O-methylbrosiperin), O-methylbalsamiferone, dihydroxanthyletin, xanthyletin, and luvangetin. *J. Chem. Soc., Perkin Trans. 1* (3), 371–376.
- Maiti, S., Panja, S.K., Bandyopadhyay, C., 2010. Substituent-controlled domino-Knoevenagel-hetero Diels-Alder reaction-a one-pot synthesis of polycyclic heterocycles. *Tetrahedron* 66 (38), 7625–7632.
- Majumdar, K.C., Bhattacharyya, T., 1997. Regioselective synthesis of furo[3,2-c][1] benzopyran-4-one and furo[3,2-c]quinolin-4-one. *J. Chem. Res. (S)* 7, 244–245.
- Mehrabi, H., Abusaidi, H., 2010. Synthesis of biscoumarin and 3,4-dihydropyrano [c] chromene derivatives catalysed by sodium dodecyl sulfate (SDS) in neat water. *J. Iran. Chem. Soc.* 7 (4), 890–894.
- Mei, R.-Q., Peng, L., Tian, F., Wang, F., Wang, L., Xu, X., Mei, R., Wang, F., 2013. Asymmetric Michael/cyclization tandem reaction of 4-hydroxycoumarin with β -nitroalkenes catalyzed by chiral bifunctional thioureas. *Org. Biomol. Chem.* 11 (8), 1286–1289.
- Metwally, M.A., Bondock, S., El-Desouky, S.I., Abdou, M.M., 2012a. A worthy insight into the dyeing applications of azo pyrazolyl dyes. *Int. J. Modern Org. Chem.* 1, 165–192.
- Metwally, M.A., Bondock, S., El-Desouky, S.I., Abdou, M.M., 2012b. Pyrazol-5-ones: tautomerism, synthesis and reactions. *Int. J. Modern Org. Chem.* 1, 19–54.
- Metwally, M.A., Bondock, S., El-Desouky, S.I., Abdou, M.M., 2012c. Synthesis, tautomeric structure, dyeing characteristics, and antimicrobial activity of novel 4-(2-arylazophenyl)-3-(2-hydroxyphenyl)-1-phenyl-2-pyrazolin-5-ones. *J. Korean Chem. Soc.* 56, 82–91.
- Metwally, M.A., Bondock, S., El-Desouky, S.I., Abdou, M.M., 2012d. Synthesis, structure investigation and dyeing assessment of novel bisazo disperse dyes derived from 3-(2-hydroxyphenyl)-1-phenyl-2-pyrazolin-5-ones for dyeing polyester fabrics. *J. Korean Chem. Soc.* 56, 348–356.
- Metwally, M.A., Bondock, S., El-Desouky, S.I., Abdou, M.M., 2013. A facile synthesis, tautomeric structure of novel 4-arylhydrazono-3-(2-hydroxyphenyl)-2-pyrazolin-5-ones and their application as disperse dyes. *Color. Technol.* 129 (6), 418–424.
- Mihelcic, D., Jakse, R., Svete, J., Stanovnik, B., Grdadolnik, S.G., 2001. Synthesis of (S)-3-heteroaryl-2-hydroxy-1-propylbenzoates by “ring switching” methodology. *J. Heterocycl. Chem.* 38 (6), 1307–1312.
- Motamedi, R., Baghbani, S., Bamoharram, F.F., 2012. Catalytic Method for synthesis of benzopyrano [3, 2-c] chromene-6, 8-dione derivatives by heteropoly acids. *Synth. Commun.* 42 (11), 1604–1612.
- Mungra, D.C., Patel, M.P., Rajani, D.P., Patel, R.G., 2011. Synthesis and identification of β -aryloxyquinolines and their pyrano[3,2-c]chromene derivatives as a new class of antimicrobial and antituberculosis agents. *Eur. J. Med. Chem.* 46 (9), 4192–4200.
- Nair, V., Menon, R.S., Vinod, A.U., Viji, S., 2002. A facile three-component reaction involving [4+1] cycloaddition leading to furan annulated heterocycles. *Tetrahedron Lett.* 43 (12), 2293–2295.
- Nair, D.K., Menna-Barreto, R.F.S., Da Silva Junior, E.N., Mobin, S.M., Namboothiri, I.N.N., 2014. Chiral squaramide-catalyzed asymmetric synthesis of pyranones and pyranonaphthoquinones via cascade reactions of 1,3-dicarbonyls with Morita-Baylis-Hillman acetates of nitroalkenes. *Chem. Commun.* 50 (53), 6973–6976.
- Nasseri, M.A., Sadeghzadeh, S.M., 2013. A highly active $\text{FeNi}_3\text{-SiO}_2$ magnetic nanoparticles catalyst for the preparation of 4H-benzo [b] pyrans and Spirooxindoles under mild conditions. *J. Iran. Chem. Soc.* 10 (5), 1047–1056.
- Naumov, M.I., Ganina, O.G., Shavirin, A.S., Beletskaya, I.P., Finet, J.P., Fedorov, A.Y., 2005. Polyfunctionalized aryllead triacetates in a cascade synthesis of tetracyclic isochromanocoumarin-type compounds. *Synthesis* 7, 1178–1182.
- Nematollahi, D., Habibi, D., Alizadeh, A., Hesari, M., 2005. An efficient conversion of catechols into 6H-benzofuro[3,2-c][1]-benzopyran-6-one derivatives. *J. Heterocycl. Chem.* 42 (2), 289–292.
- Nesterov, V.N., Wiedenfeld, D., Nesterova, S.V., Daniels, L.M., 2005. Synthesis and X-ray structural investigations of 2-amino-7-methyl-4-(1-naphthyl)-5-oxo-4H,5H-pyrano [4,3-b]pyran-3-carbonitrile and 2-amino-4-(1-naphthyl)-5-oxo-4H,5H-pyrano [3,2-c]chromene-3-carbonitrile. *J. Chem. Crystallogr.* 35 (11), 917–922.
- Niknam, K., Jamali, A., 2012. Silica-bonded N-propylpiperazine sodium n-propionate as recyclable basic catalyst for synthesis of 3,4-dihydropyrano[c]chromene derivatives and biscoumarins. *Chin. J. Catal.* 33 (11), 1840–1849.
- Nishino, H., Kumabe, R., Hamada, R., Yakut, M., 2014. Mn(III)-based reaction of alkenes with quinolinones. Formation of peroxyquinolinones and quinoline-related derivatives. *Tetrahedron* 70 (7), 1437–1450.
- Okamoto, Y., Okawara, T., Furukawa, M.A., 1996. Short three components approach to fused pyrrolo[2,3-b]pyrrolidinone. *Heterocycles* 43 (11), 2353–2356.
- Okamoto, Y., Kaneda, Y., Yamasaki, T., Okawara, T., Furukawa, M., 1997. Simple preparation of fused pyrrolo[2,3-b]pyrrolidinones and pyrrolo[2,3-c]pyridazinones. *J. Chem. Soc., Perkin Trans. 1* (9), 1323–1327.
- Pal, S., Khan, M.N., Karamthulla, S., Choudhury, L.H., 2013. Molecular iodine catalyzed one-pot multicomponent reactions for the synthesis of dihydrochromeno [4, 3-b] pyrazolo [4, 3-e] pyridin-6 (7 H)-ones. *RSC Adv.* 3 (36), 15705–15711.
- Pandya, S.U., Pandya, U.R., Hirani, B.R., Brahmabhatt, D.I., 2006. One-pot synthesis of diarylpyrido[3,2-c]coumarins. *J. Heterocycl. Chem.* 43 (3), 795–798.
- Panja, S.K., Ghosh, J., Maiti, S., Bandyopadhyay, C., 2012. Synthesis of furocoumarins and biscoumarins by an isocyanide-induced multicomponent reaction: isocyanide as a masked amine. *J. Chem. Res.* 36 (4), 222–225.
- Pansuriya, A.M., Savant, M.M., Bhuva, C.V., Singh, J., Naliapara, Y.T., 2009. Use of cyclic aliphatic ketones for spiro 2-amino-3-cyano pyrano [3, 2-c] chromene formation. *ARKIVOC* xii, 254–260.
- Ponpandian, T., Muthusubramanian, S., 2014. One-pot, catalyst-free synthesis of spirooxindole and 4 h-pyran derivatives. *Synth. Commun.* 44 (6), 868–874.
- Prasanna, T.S.R., Raju, K.M., 2011. Silica gel promoted mild, efficient and inexpensive protocol for the preparation of 3,4-dihydropyrano[c]chromenes. *J. Korean Chem.* 55 (4), 662–665.
- Rad-Moghadam, K., Taghizadeh, V., Alireza, Alipour, A., 2014. LiBF_4 integrated into $[\text{BMI}_m]\text{BF}_4$: an ionic-liquid metal composite and homogeneous catalyst for efficient synthesis of pyran-annulated heterocycles. *Appl. Organomet. Chem.* 28 (3), 146–150.
- Rajawat, A., Khandelwal, S., Kumar, M., 2014. Deep eutectic solvent promoted efficient and environmentally benign four-component domino protocol for synthesis of spirooxindoles. *RSC Adv.* 4 (10), 5105–5112.
- Ray, S., Paul, S.K., 2004. Studies on reactions of hydroxycoumarin compounds with formamide. *J. Indian Chem. Soc.* 81 (6), 488–491.
- Sabitha, G., Reddy, G.S.K.K., Reddy, K.B., Yadav, J.S., 2003. Vanadium (III) chloride catalyzed Biginelli condensation: Solution phase generation of dihydropyrimidin-(2H)-ones. *Tetrahedron Lett.* 44 (34), 6497–6499.

- Saha, P., 2013. Fermented baker's yeast: an efficient catalyst for the synthesis of pyran derivatives in water at room temperature. *Synth. Commun.* 43 (12), 1708–1713.
- Sarma, R., Sarmah, M.M., Lekhok, K.C., Prajapati, D., 2010. Organic reactions in water: an efficient synthesis of pyranocoumarin derivatives. *Synlett* 19, 2847–2852.
- Sato, K., Yokoo, E., Takenaga, N., 2013. Facile synthesis of guaiazulene-heterocycle hybrids via multicomponent reactions involving formation of zwitterionic intermediates. *Heterocycles* 87 (4), 807–814.
- Seifi, M., Sheibani, H., 2008. High surface area MgO as a highly effective heterogeneous base catalyst for three-component synthesis of tetrahydrobenzopyran and 3,4-dihydropyrano[c]chromene derivatives in aqueous media. *Catal. Lett.* 126 (3–4), 275–279.
- Shaabani, A., Teimouri, M.B., 2003. Microwave-assisted rapid synthesis of furan annulated heterocycles. *J. Chem. Res. (S)* 11, 732–733.
- Shaabani, A., Teimouri, M.B., Bijanzadeh, H.R., 2004. One-pot three-component condensation reactions in water. An efficient and improved procedure for the synthesis of furan annulated heterocycles. *Monatsh. Chem.* 135 (5), 589–593.
- Shaabani, A., Samadi, S., Badri, Z., Rahmati, A., 2005a. Ionic liquid promoted efficient and rapid one-pot synthesis of pyran annulated heterocyclic systems. *Catal. Lett.* 104 (1–2), 39–43.
- Shaabani, A., Teimouri, M.B., Samadi, S., Soleimani, K., 2005b. Microwave-assisted three-component condensation on montmorillonite K10. Solvent-free synthesis of furopyrimidines, furocoumarins, and furopyranones. *Synth. Commun.* 35 (4), 535–541.
- Shaabani, A., Samadi, S., Rahmati, A., 2007. One-pot, three-component condensation reaction in water: an efficient and improved procedure for the synthesis of pyran annulated heterocyclic systems. *Synth. Commun.* 37 (3), 491–499.
- Shaabani, A., Rezayan, A.H., Sarvary, A., Rahmati, A., 2008. Isocyanide-catalyzed reaction of tetracyanoethylene and activated 1,3-dicarbonyl CH-acid compounds. A rapid and efficient synthesis of pyran annulated heterocyclic systems. *Synth. Commun.* 38 (2), 274–281.
- Shafiee, A., Motamedi, R., Firuzi, O., Meili, S., Mehdipour, A.R., Miri, R., 2011. Synthesis and cytotoxic activity of novel benzopyrano [3, 2-c] chromene-6, 8-dione derivatives. *Med. Chem. Res.* 20 (4), 466–474.
- Shaterian, H.R., Oveisi, A.R.A., 2011. Simple green approach to the synthesis of 2-amino-5-oxo-4,5-dihydropyrano[3,2-c]chromene-3-carbonitrile derivatives catalyzed by 3-hydroxypropanaminium acetate (HPAA) as a new ionic liquid. *J. Iran. Chem. Soc.* 8 (2), 545–552.
- Shestopalov, A.A., Rodinovskaya, L.A., Shestopalov, A.M., Litvinov, V.P., 2005. Cross reactions of cyanoacetic acid derivatives with carbonyl compounds. One-step synthesis of substituted 2-amino-5-oxo-4,5-dihydropyrano[3,2-c]chromenes. *Russ. Chem. Bull.* 54 (4), 992–996.
- Siddiqui, Z.N., 2014. Sulfamic acid catalysed synthesis of pyranocoumarins in aqueous media. *Tetrahedron Lett.* 55 (1), 163–168.
- Siddiqui, Z.N., Khan, T., 2013. P_2O_5/SiO_2 as an efficient heterogeneous catalyst for the synthesis of heterocyclic alkene derivatives under thermal solvent-free conditions. *Catal. Sci. Technol.* 3 (8), 2032–2043.
- Siddiqui, Z.N., Khan, T., 2014. Unprecedented single-pot protocol for the synthesis of novel bis-3, 4-dihydropyrimidin-2 (1H)-ones using PEG-HClO₄ as a biodegradable, highly robust and reusable solid acid green catalyst under solvent-free conditions. *RSC Adv.* 4 (5), 2526–2537.
- Sorsak, G., Grdadolnik, S.G., Stanovnik, B., 1998. The synthesis of ethyl 2-[(2,2-dibenzoyl) ethenyl]amino-3-dimethylaminopropenoate and its application to the synthesis of fused 3-aminopyran-2-ones and 3-aminoazolo- and -azinopyrimidin-4(4H)-ones. *J. Heterocycl. Chem.* 35 (6), 1275–1279.
- Strakova, I., Strakovs, A., Petrova, M., Belyakov, S., 2009. Synthesis and reactions of 4-(arylhdyrazino) coumarins. *Chem. Heterocycl. Compd.* 45 (11), 1319–1324.
- Sun, X.J., Zhou, J.F., Zhi, S.J., 2012. Efficient one-pot synthesis of tetrahydrobenzo [c] xanthene-1, 11-dione derivatives under microwave irradiation. *Synth. Commun.* 42 (13), 1987–1994.
- Tabatabaiean, K., Heidari, H., Mamaghani, M., Mahmoodi, N.O., 2012. Ru (II) complexes bearing tertiary phosphine ligands: a novel and efficient homogeneous catalyst for one-pot synthesis of dihydropyrano [3, 2-c] chromene and tetrahydrobenzo [b] pyran derivatives. *Appl. Organomet. Chem.* 26 (2), 56–61.
- Tietze, L.F., Evers, H., Topken, E., 2001. A novel concept in combinatorial chemistry in solution with the advantages of solid-phase synthesis: formation of N-betaines by multicomponent domino reactions. *Angew. Chem. Int. Ed.* 40 (5), 903–905.
- Toche, R.B., Jachak, M.N., Sabnis, R.W., Kappe, T., 1999. Synthesis of novel pyran fused quinolones, coumarins, and pyridines. *J. Heterocycl. Chem.* 36 (2), 467–471.
- Toplak, R., Selic, L., Sorsak, G., Stanovnik, B., 1997. Reactions of ethyl (Z)-2-[2,2-bis(ethoxycarbonyl) vinyl]amino-3-dimethylaminopropenoate with C-nucleophiles. Synthesis of substituted 3-amino-2H-pyran-2-ones. *Heterocycles* 45 (3), 555–565.
- Toplak, R., Svete, J., Stanovnik, B., Grdadolnik, S.G., 1999. The synthesis of methyl 2-(benzyloxycarbonyl)amino-3-dimethylaminopropenoate. The synthesis of trisubstituted pyrroles, 3-amino-2H-pyran-2-ones, fused 2H-pyran-2-ones and 4H-pyridin-4-ones. *J. Heterocycl. Chem.* 36 (1), 225–235.
- Tu, S., Jiang, H., Fang, F., Feng, Y., Zhu, S., Li, T., Zhang, X., Shi, D., 2004. Synthesis of 2-amino-3-ethoxycarbonyl-4-aryl-4H,5H-pyrano-[3,2-c]benzopyran-5-one. *J. Chem. Res.* 6, 396–398.
- Vovk, M.V., Sukach, V.A., Dorokhov, V.I., 2007. Heterocyclization of functionalized heterocumulenes with C, N-, C, O-, and C, S-bimucleophiles. Part VIII. Synthesis of pyrano(chromeno)[3,4-e][1,3]oxazines by condensation of 1-chloroalkyl isocyanates with 4-hydroxy-6-methylpyran-2-one and 4-hydroxycoumarin. *Russ. J. Org. Chem.* 43 (8), 1186–1189.
- Wang, J., Shi, D., Zhuang, Q., Wang, X., Hu, H., 2004. Synthesis of 4H,5H-pyrano[3,2-c][1] benzopyran-5-ones in aqueous media. *J. Chem. Res.* 12, 818–820.
- Wang, X.S., Zhang, M.M., Zeng, Z.S., Shi, D.Q., Tu, S.J., Wei, X.Y., Zong, Z.M., 2005. A clean procedure for the synthesis of chromeno[4,3-b]benzo[f]quinoline and quinolino[4,3-b]benzo[f]quinoline derivatives in aqueous media. *Chem. Lett.* 34 (10), 1316–1317.
- Wang, X.S., Zhang, M.M., Zeng, Z.S., Shi, D.Q., Tu, S.J., 2006. Clean procedure for synthesis of chromeno[4,3-b]benzo[f]quinolin-6-one derivatives. Reaction of N-arylideneaphthalen-2-amine with 4-hydroxycoumarin in aqueous media. *Synth. Commun.* 36 (14), 2047–2057.
- Wang, H.J., Lu, J., Zhang, Z.H., 2010. Highly efficient three-component, one-pot synthesis of dihydropyrano [3, 2-c] chromene derivatives. *Monatsh. Chem.* 141 (10), 1107–1112.
- Wu, J., 2006. General microwave-assisted protocols for the expedient synthesis of furo[3,2-c]chromen-4-one. *Chem. Lett.* 35 (1), 118–119.
- Xiao, J., Chen, Y., Zhu, S., Wang, L., Xu, L., Wei, H., 2014. Diversified construction of chromeno[3,4-c]pyridin-5-one and benzo[c]chromen-6-one derivatives by domino reaction of 4-Alkynyl-2-oxo-2H-chromene-3-carbaldehydes. *Adv. Synth. Catal.* 356 (8), 1835–1845.
- Xie, J.W., Li, P., Wang, T., Zhou, F.T., 2011. Efficient and mild synthesis of functionalized 2,3-dihydrofuran derivatives via domino reaction in water. *Tetrahedron Lett.* 52 (18), 2379–2382.
- Yao, M.J., Guan, Z., He, Y.H., 2013. Simple, catalyst-free, one-pot procedure for the synthesis of 2-amino-3-cyano-1, 4, 5, 6-tetrahydropyrano [3, 2-c] quinolin-5-one derivatives. *Synth. Commun.* 43 (15), 2073–2078.

- Yavari, I., Sabbaghan, M., Hossaini, Z., 2008. Proline-promoted efficient synthesis of 4-aryl-3,4-dihydro-2H,5H-pyrano[3,2-c]chromene-2,5-diones in aqueous media. *Synlett* 8, 1153–1154.
- Yilmaz, M., Yakut, M., Pekel, A.T., 2008. Synthesis of 2,3-dihydro-4H-furo[3,2-c]chromen-4-ones and 2,3-dihydronaphtho[2,3-b]furan-4,9-diones by the radical cyclizations of hydroxyenones with electron-rich alkenes using manganese(III) acetate. *Synth. Commun.* 38 (6), 914–927.
- Zhao, B.L., Du, D.M., 2014. Enantioselective synthesis of enol lactones from tandem Michael addition/lactonization catalyzed by a chiral squaramide catalyst. *Tetrahedron Asymmetry* 25 (4), 310–317.
- Zhao, P.L., Wang, L., Zhu, X.L., Huang, X., Zhan, C.G., Wu, J.W., Yang, G.F., 2010. Subnanomolar inhibitor of cytochrome bcl complex designed by optimizing interaction with conformationally flexible residues. *J. Am. Chem. Soc.* 132 (1), 185–194.
- Zheng, J., Li, Y.Q., 2011. One-pot synthesis of tetrahydrobenzo [b] pyran and dihydropyrano [c] chromene derivatives in aqueous media by using trisodium citrate as a green catalyst. *Arch. Appl. Sci. Res.* 3 (2), 38–388.
- Zhu, X., Xu, X.P., Sun, C., Chen, T., Shen, Z.L., Ji, S.J., 2011. I-MCR-Ullmann cascade toward furo [2,3-b]indole scaffold. *Tetrahedron* 67 (34), 6375–6381.
- Zhuang, Q., Zhou, D., Tu, S., Li, C., Cao, L., Shao, Q., 2008. A highly efficient microwave-assisted synthesis of chromeno[3,4-b][4,7]phenanthroline derivatives through multicomponent reactions in water. *J. Heterocycl. Chem.* 45 (3), 831–835.
- Ziarani, G.M., Hajiabbasi, P., 2013. Recent application of 4-hydroxycoumarin in multi-component reactions. *Heterocycles* 87 (7), 1415–1440.